

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

TIM3 / HAVCR2/ CD366 (Effector T-Cell Marker) Antibody, IgG2b, Clone: [TIM3/4025], Mouse, Monoclonal NBT-84868-MSM5-P0

Article Name	TIM3 / HAVCR2/ CD366 (Effector T-Cell Marker) Antibody, IgG2b, Clone: [TIM3/4025], Mouse, Monoclonal
Biozol Catalog Number	NBT-84868-MSM5-P0
Supplier Catalog Number	84868-MSM5-P0
Alternative Catalog Number	NBT-84868-MSM5-P0-20,NBT-84868-MSM5-P0-100
Manufacturer	NeoBiotechnologies
Host	Mouse
Category	Antikörper
Application	IHC, WB
Species Reactivity	Human
Immunogen	Recombinant fragment of human TIM3 protein (around aa 22-202) (exact sequence is proprietary)
Product Description	TIMs are type I transmembrane glycoproteins with one Ig-like V-type domain and a Ser/Thr-rich mucin stalk. TIM-3 is expressed on the surface of effector T cel tonsil, lymph node or spleen. Is (CD4+Th1 and CD8+Tc1) but not on helper T cells (CD4+Th2 a
Clonality	Monoclonal
Clone Designation	[TIM3/4025]
Molecular Weight	30-60kDa
Isotype	lgG2b

NCBI	84868
UniProt	Q8TDQ0
Form	200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with0.05% BSA & D.05% azide. Also available WITHOUT BSA & D.05% azide at 1.0mg/ml.
Antibody Type	Monoclonal Antibody
Application Notes	Western Blot (1-2ug/ml),,Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 2