

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

TIM3 / HAVCR2/ CD366 (Effector T-Cell Marker) Antibody, IgG2b, Clone: [TIM3/4027], Mouse, Monoclonal NBT-84868-MSM7-P0

| | |
|----------------------------|---|
| Article Name | TIM3 / HAVCR2/ CD366 (Effector T-Cell Marker) Antibody, IgG2b, Clone: [TIM3/4027], Mouse, Monoclonal |
| Biozol Catalog Number | NBT-84868-MSM7-P0 |
| Supplier Catalog Number | 84868-MSM7-P0 |
| Alternative Catalog Number | NBT-84868-MSM7-P0-20,NBT-84868-MSM7-P0-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC, WB |
| Species Reactivity | Human |
| Immunogen | Recombinant fragment of human TIM3 protein (around aa 22-202) (exact sequence is proprietary) |
| Product Description | TIMs are type I transmembrane glycoproteins with one Ig-like V-type domain and a Ser/Thr-rich mucin stalk. TIM-3 is expressed on the surface of effector T cell tonsil, lymph node or spleen. Is (CD4+Th1 and CD8+Tc1) but not on helper T cells (CD4+Th2 ... |
| Clonality | Monoclonal |
| Clone Designation | [TIM3/4027] |
| Molecular Weight | 60kDa |
| Isotype | IgG2b |

| | |
|-------------------|---|
| NCBI | 84868 |
| UniProt | Q8TDQ0 |
| Form | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Application Notes | Western Blot (1-2ug/ml), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT), (Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 2 |