

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Prostein (p501S) / Solute Carrier Family 45 Member 3 (SLC45A3) Antibody, IgG1, Clone: [SLC45A3/7647], Mouse, Monoclonal NBT-85414-MSM7-P0

| Article Name               | Prostein (p501S) / Solute Carrier Family 45 Member 3 (SLC45A3)<br>Antibody, IgG1, Clone: [SLC45A3/7647], Mouse, Monoclonal  |
|----------------------------|---|
| Biozol Catalog Number      | NBT-85414-MSM7-P0   |
| Supplier Catalog Number    | 85414-MSM7-P0   |
| Alternative Catalog Number | NBT-85414-MSM7-P0-20,NBT-85414-MSM7-P0-100  |
| Manufacturer               | NeoBiotechnologies  |
| Host                       | Mouse   |
| Category                   | Antikörper  |
| Application                | IHC   |
| Species Reactivity         | Human   |
| Immunogen                  | Recombinant human SLC45A3 protein   |
| Product Description        | Human prostein is a 553 aa protein identified by cDNA library subtraction and subsequent high throughput microarray screening of prostate cancer. Prostein has multiple transmembrane domains. Prostein has been shown to be uniquely expressed in normal |
| Clonality                  | Monoclonal  |
| Clone Designation          | [SLC45A3/7647]  |
| Molecular Weight           | 59kDa   |
| Isotype                    | IgG1  |

| NCBI              | 85414  |
|-------------------|--|
| UniProt           | Q96JT2   |
| Form              | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are a side. Also available WITHOUT BSA & Double at 1.0mg/ml.  |
| Antibody Type     | Monoclonal Antibody  |
| Application Notes | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95&degC followed by cooling at RT for 20 minutes),Optimal dilutio |