

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Recombinant p40 (deltaNp63) (Squamous, Basal & Myoepithelial Cell Marker) Antibody, IgG1, Clone: [rTP40/3690], Mouse, Monoclonal NBT-8626-MSM10-P0

| | |
|----------------------------|---|
| Article Name | Recombinant p40 (deltaNp63) (Squamous, Basal & Myoepithelial Cell Marker) Antibody, IgG1, Clone: [rTP40/3690], Mouse, Monoclonal |
| Biozol Catalog Number | NBT-8626-MSM10-P0 |
| Supplier Catalog Number | 8626-MSM10-P0 |
| Alternative Catalog Number | NBT-8626-MSM10-P0-20,NBT-8626-MSM10-P0-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Mouse |
| Category | Antikörper |
| Application | FC, IF, IHC |
| Species Reactivity | Human |
| Immunogen | A synthetic peptide from the N-terminal of human p40protein (Exact sequence is proprietary) |
| Product Description | p63 consists of two major isoforms-TAp63 and delta-Np63. These isoforms differ in the structure of the N-terminal domains. The TAp63 isoform (identified by anti-p63 antibody) contains a transactivation-competent TA domain with homology to p53, which ... |
| Clonality | Monoclonal |
| Clone Designation | [rTP40/3690] |
| Molecular Weight | 40kDa |
| Isotype | IgG1 |

| | |
|-------------------|---|
| NCBI | 8626 |
| UniProt | Q9H3D4 |
| Form | 200ug/ml of Ab purified by Protein A Column. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Recombinant Monoclonal Antibody |
| Application Notes | Flow Cytometry (1-2ug/million cells), Immunofluorescence (1-2ug/ml), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 mi |