

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Recombinant p40 (Squamous, Basal & Myoepithelial Cell Marker) Antibody, Clone: [P40/4396R], Rabbit, Monoclonal NBT-8626-RBM14-P0

| Article Name               | Recombinant p40 (Squamous, Basal & Myoepithelial Cell Marker)<br>Antibody, Clone: [P40/4396R], Rabbit, Monoclonal  |
|----------------------------|--|
| Biozol Catalog Number      | NBT-8626-RBM14-P0  |
| Supplier Catalog Number    | 8626-RBM14-P0  |
| Alternative Catalog Number | NBT-8626-RBM14-P0-20,NBT-8626-RBM14-P0-100   |
| Manufacturer               | NeoBiotechnologies   |
| Host                       | Rabbit   |
| Category                   | Antikörper   |
| Application                | ELISA, FC, IHC   |
| Species Reactivity         | Human  |
| Immunogen                  | Recombinant fragment (around aa1-100) of human p40 protein (exact sequence is proprietary)   |
| Product Description        | p40 antibody (antibody recognizing Np63 only) is an isoform of p63, is thought to function as a stem cell factor, responsible for maintaining cells in an uncommitted state with regenerative potential a role that may be recapitulated in tumors derived |
| Clonality                  | Monoclonal   |
| Clone Designation          | [P40/4396R]  |
| Molecular Weight           | 63kDa  |
| NCBI                       | 8626   |

| UniProt           | Q9H3D4   |
|-------------------|--|
| Form              | 200ug/ml of Ab purified by Protein A Column. Prepared in 10mM PBS with 0.05% BSA & D.05% azide. Also available WITHOUT BSA & D.05% azide at 1.0mg/ml.  |
| Antibody Type     | Recombinant Monoclonal Antibody  |
| Application Notes | ELISA (For coating, order antibody without BSA), Flow Cytometry (1-2ug/million cells), ,Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, |