

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

CD2 / Lymphocyte Function Antigen 2 (LFA-2) Antibody, IgG1, Clone: [LFA2/7106], Mouse, Monoclonal NBT-914-MSM16-P0

| | |
|----------------------------|---|
| Article Name | CD2 / Lymphocyte Function Antigen 2 (LFA-2) Antibody, IgG1, Clone: [LFA2/7106], Mouse, Monoclonal |
| Biozol Catalog Number | NBT-914-MSM16-P0 |
| Supplier Catalog Number | 914-MSM16-P0 |
| Alternative Catalog Number | NBT-914-MSM16-P0-20,NBT-914-MSM16-P0-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC, WB |
| Species Reactivity | Human |
| Immunogen | Recombinant fragment of human CD2 protein (exact sequence is proprietary) |
| Product Description | CD2 is mainly present on the surface of mature T cells and NK cell membranes, while B cells are usually not expressed. It can be used to label the diagnosis and study of normal T cells and their associated tumors, such as pre-t cell lymphoma, margina... |
| Clonality | Monoclonal |
| Clone Designation | [LFA2/7106] |
| Molecular Weight | 39kDa |
| Isotype | IgG1 |

| | |
|-------------------|---|
| NCBI | 914 |
| UniProt | P06729 |
| Form | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Application Notes | Western Blot (1-2ug/ml), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT), (Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 2 |