

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Recombinant CD2 / Lymphocyte Function Antigen 2 (LFA-2) Antibody, IgG1, Clone: [rLFA2/7147], Mouse, Monoclonal NBT-914-MSM17-P1ABX

| | |
|----------------------------|---|
| Article Name | Recombinant CD2 / Lymphocyte Function Antigen 2 (LFA-2) Antibody, IgG1, Clone: [rLFA2/7147], Mouse, Monoclonal |
| Biozol Catalog Number | NBT-914-MSM17-P1ABX |
| Supplier Catalog Number | 914-MSM17-P1ABX |
| Alternative Catalog Number | NBT-914-MSM17-P1ABX-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human |
| Immunogen | Recombinant fragment of human CD2 protein corresponding to N-terminal (exact sequence is proprietary) |
| Product Description | CD2 is mainly present on the surface of mature T cells and NK cell membranes, while B cells are usually not expressed. It can be used to label the diagnosis and study of normal T cells and their associated tumors, such as pre-t cell lymphoma, margina... |
| Clonality | Monoclonal |
| Clone Designation | [rLFA2/7147] |
| Molecular Weight | 39kDa |
| Isotype | IgG1 |

| | |
|-------------------|--|
| NCBI | 914 |
| UniProt | P06729 |
| Form | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Recombinant Monoclonal Antibody |
| Application Notes | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 20 minutes),Optimal diluti |