

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

CD6 (Negative Marker of T-regulatory Cells) Antibody, IgG1, Clone: [3F7B5], Mouse, Monoclonal NBT-923-MSM1-P0

| Article Name | CD6 (Negative Marker of T-regulatory Cells) Antibody, IgG1, Clone: [3F7B5], Mouse, Monoclonal |
|----------------------------|---|
| Biozol Catalog Number | NBT-923-MSM1-P0 |
| Supplier Catalog Number | 923-MSM1-P0 |
| Alternative Catalog Number | NBT-923-MSM1-P0-20,NBT-923-MSM1-P0-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Mouse |
| Category | Antikörper |
| Application | FC, IF, IHC |
| Species Reactivity | Human |
| Immunogen | Human rheumatoid synovial T cell line ST-1 |
| Product Description | CD6 is a type I transmembrane glycoprotein that contains a 24- amino acid signal sequence, three extracellular scavenger receptor cysteine-rich (SRCR) domains, a membrane-spanning domain and a 44-amino acid cytoplasmic domain. The CD6 glycoprotein is |
| Clonality | Monoclonal |
| Clone Designation | [3F7B5] |
| Molecular Weight | 90-130kDa |
| Isotype | lgG1 |

| NCBI | 923 |
|-------------------|--|
| UniProt | P30203 |
| Form | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & amp; 0.05% azide. Also available WITHOUT BSA & amp; azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Application Notes | Flow Cytometry (1-2ug/million cells), Immunofluorescence (1- 2ug/ml), ,Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 mi |