

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **CD20 / MS4A1 (B-Cell Marker) Antibody, IgG2a, Clone: [L26], Mouse, Monoclonal NBT-931-MSM2-P1ABX**

|                            |   |
|----------------------------|---|
| Article Name               | CD20 / MS4A1 (B-Cell Marker) Antibody, IgG2a, Clone: [L26], Mouse, Monoclonal   |
| Biozol Catalog Number      | NBT-931-MSM2-P1ABX  |
| Supplier Catalog Number    | 931-MSM2-P1ABX  |
| Alternative Catalog Number | NBT-931-MSM2-P1ABX-100  |
| Manufacturer               | NeoBiotechnologies  |
| Host                       | Mouse   |
| Category                   | Antikörper  |
| Application                | FC, IF, WB  |
| Species Reactivity         | Human   |
| Immunogen                  | Human tonsil B cells  |
| Product Description        | Recognizes a protein of 30-33kDa, which is identified as CD20. Its epitope is located in the cytoplasmic domain of CD20 and was, therefore, ascribed as CD20cy in the 5th Workshop. CD20 is a non-Ig differentiation antigen of B-cells and its expression... |
| Clonality                  | Monoclonal  |
| Clone Designation          | [L26]   |
| Molecular Weight           | 33-37kDa  |
| Isotype                    | IgG2a   |

|                   |   |
|-------------------|---|
| NCBI              | <a href="#">931</a>   |
| UniProt           | <a href="#">P11836</a>  |
| Form              | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.  |
| Antibody Type     | Monoclonal Antibody   |
| Application Notes | Flow Cytometry (0.5-1ug/million cells), Immunofluorescence (1-2ug/ml), Western Blotting (1-2ug/ml), Immunohistology (Formalin-fixed) (0.5-1ug/ml for 30 minutes at RT), (Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate bu |