

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Recombinant CD22 / BL-CAM (B-Cell Marker) Antibody, Clone: [BLCAM/8113R], Rabbit, Monoclonal NBT-933-RBM10-P1ABX

| Article Name | Recombinant CD22 / BL-CAM (B-Cell Marker) Antibody, Clone: [BLCAM/8113R], Rabbit, Monoclonal |
|----------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Biozol Catalog Number | NBT-933-RBM10-P1ABX |
| Supplier Catalog Number | 933-RBM10-P1ABX |
| Alternative Catalog Number | NBT-933-RBM10-P1ABX-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Rabbit |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human |
| Immunogen | Recombinant fragment (around aa1-200) of human BL-CAM/CD22 protein (exact sequence is proprietary) |
| Product Description | Recognizes a protein of 130-140kDa, identified as CD22 (also known as BL-CAM). CD22 expression is restricted to normal and neoplastic B cells and is absent from other haemopoietic cell types. In B-cell ontogeny, CD22 is first expressed in the cytopla |
| Clonality | Monoclonal |
| Clone Designation | [BLCAM/8113R] |
| Molecular Weight | 130-140kDa |
| NCBI | 933 |

| UniProt | P20273 |
|-------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Form | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & amp; 0.05% azide. Also available WITHOUT BSA & amp; azide at 1.0mg/ml. |
| Antibody Type | Recombinant Monoclonal Antibody |
| Application Notes | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes),Optimal dilution f |