

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Recombinant MUC16 / CA125 (Ovarian Carcinoma Marker) Antibody, Clone: [OCA125/4505R], Rabbit, Monoclonal NBT-94025-RBM5-P0

| Article Name | Recombinant MUC16 / CA125 (Ovarian Carcinoma Marker) Antibody, Clone: [OCA125/4505R], Rabbit, Monoclonal |
|----------------------------|---|
| Biozol Catalog Number | NBT-94025-RBM5-P0 |
| Supplier Catalog Number | 94025-RBM5-P0 |
| Alternative Catalog Number | NBT-94025-RBM5-P0-20,NBT-94025-RBM5-P0-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Rabbit |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human |
| Immunogen | Synthetic peptide corresponding to CA125 residues within aa12201 12300 of CA125 was used as an immunogen. |
| Product Description | CA125 also known as mucin 16, is a member of the mucin family glycoproteins. CA125 is a component of the ocular surface (including the cornea and conjunctiva), the respiratory tract and the female reproductive tract epithelia. Since CA125 is highly g |
| Clonality | Monoclonal |
| Clone Designation | [OCA125/4505R] |
| Molecular Weight | >2 000kDa |
| NCBI | 94025 |

| UniProt | Q8WXI7 |
|-------------------|--|
| Form | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & amp; 0.05% azide. Also available WITHOUT BSA & amp; azide at 1.0mg/ml. |
| Antibody Type | Recombinant Monoclonal Antibody |
| Application Notes | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 20 minutes),Optimal diluti |