

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **CD30 / TNFRSF8 (Hodgkin & Reed-Sternberg Cell Marker) Antibody, IgG1, Clone: [Ber-H2], Mouse, Monoclonal NBT-943-MSM2-CS0**

Article Name	CD30 / TNFRSF8 (Hodgkin & Reed-Sternberg Cell Marker) Antibody, IgG1, Clone: [Ber-H2], Mouse, Monoclonal
Biozol Catalog Number	NBT-943-MSM2-CS0
Supplier Catalog Number	943-MSM2-CS0
Alternative Catalog Number	NBT-943-MSM2-CS0-0.1,NBT-943-MSM2-CS0-0.5,NBT-943-MSM2-CS0-100
Manufacturer	NeoBiotechnologies
Host	Mouse
Category	Antikörper
Application	IHC
Species Reactivity	Human
Immunogen	Co cell line established from a patient with Hodgkins disease of T-cell lineage
Product Description	Recognizes a single chain glycoprotein of 105/120kDa, identified as CD30/Ki-1. Its epitope is located between aa112-412. CD30 is synthesized as a 90kDa precursor, which is processed in the Golgi complex into a membrane-bound phosphorylated mature 105...
Clonality	Monoclonal
Clone Designation	[Ber-H2]
Molecular Weight	105-120kDa

Isotype	IgG1
NCBI	<a href="#">943</a>
UniProt	<a href="#">P28908</a>
Form	Tissue culture supernatant with 0.05% Azide. Contact us if you require it in a different format.
Antibody Type	Monoclonal Antibody
Application Notes	Immunohistochemistry (Formalin-fixed) (1:50-1:100 for 30 minutes at RT) (Staining of formalin-fixed tissues requires heating tissue sections in 1mM EDTA, pH 7.5-8.5, for 45 min at 95&degC followed by cooling at RT for 20 minutes)Optimal dilution for a sp