

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

CD30 / TNFRSF8 (Hodgkin & Reed-Sternberg Cell Marker) Antibody, IgG1, Clone: [SPM121], Mouse, Monoclonal NBT-943-MSM2X-CS0

CD30 / TNFRSF8 (Hodgkin & Reed-Sternberg Cell Marker) Antibody, IgG1, Clone: [SPM121], Mouse, Monoclonal
NBT-943-MSM2X-CS0
943-MSM2X-CS0
NBT-943-MSM2X-CS0-0.1,NBT-943-MSM2X-CS0-0.5,NBT-943-MSM2X-CS0-100
NeoBiotechnologies
Mouse
Antikörper
IHC
Human
Co cell line established from a patient with Hodgkins disease of T-cell lineage
Recognizes a single chain glycoprotein of 105/120kDa, identified as CD30/Ki-1. CD30 is synthesized as a 90kDa precursor, which is processed in the Golgi complex into a membrane-bound phosphorylated mature 105/120kDa glycoprotein. In Hodgkins disease,
Monoclonal
[SPM121]
105-120kDa

Isotype	lgG1
NCBI	943
UniProt	P28908
Form	Tissue culture supernatant with 0.05% Azide. Contact us if you require it in a different format.
Antibody Type	Monoclonal Antibody
Application Notes	Immunohistochemistry (Formalin-fixed) (1:50-1:100 for 30 minutes at RT) (Staining of formalin-fixed tissues requires heating tissue sections in 1mM EDTA, pH 7.5-8.5, for 45 min at 95°C followed by cooling at RT for 20 minutes)Optimal dilution for a sp