

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **Napsin A (Lung Adenocarcinoma Marker) Antibody, IgG1, Clone: [NAPSA/3308], Mouse, Monoclonal NBT-9476-MSM10-P0**

|                            |  |
|----------------------------|--|
| Article Name               | Napsin A (Lung Adenocarcinoma Marker) Antibody, IgG1, Clone: [NAPSA/3308], Mouse, Monoclonal   |
| Biozol Catalog Number      | NBT-9476-MSM10-P0  |
| Supplier Catalog Number    | 9476-MSM10-P0  |
| Alternative Catalog Number | NBT-9476-MSM10-P0-20,NBT-9476-MSM10-P0-100   |
| Manufacturer               | NeoBiotechnologies   |
| Host                       | Mouse  |
| Category                   | Antikörper   |
| Application                | IHC  |
| Species Reactivity         | Human  |
| Immunogen                  | Recombinant human Napsin A protein fragment (aa189-299) (exact sequence is proprietary)  |
| Product Description        | Napsin is a pepsin-like aspartic proteinase connected with maturation of surfactant protein B. There are two closely related napsins, napsin A and napsin B. Napsin A is expressed as a single chain protein. Immunohistochemical studies revealed high exp... |
| Clonality                  | Monoclonal   |
| Clone Designation          | [NAPSA/3308]   |
| Molecular Weight           | 37kDa  |
| Isotype                    | IgG1   |

|                   |  |
|-------------------|--|
| NCBI              | <a href="#">9476</a>   |
| UniProt           | <a href="#">O96009</a>   |
| Form              | 200ug/ml of Ab Purified from rabbit anti-serum by Protein A. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA at 1.0mg/ml.  |
| Antibody Type     | Monoclonal Antibody  |
| Application Notes | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes), Optimal dilution |