

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Lewis A (Blood Group Antigen) Antibody, IgG1, Clone: [7LE], Mouse, Monoclonal NBT-MSM1-311-P0

| | |
|----------------------------|---|
| Article Name | Lewis A (Blood Group Antigen) Antibody, IgG1, Clone: [7LE], Mouse, Monoclonal |
| Biozol Catalog Number | NBT-MSM1-311-P0 |
| Supplier Catalog Number | MSM1-311-P0 |
| Alternative Catalog Number | NBT-MSM1-311-P0-20,NBT-MSM1-311-P0-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Mouse |
| Category | Antikörper |
| Application | FC, IF, IHC |
| Species Reactivity | Human, Mouse |
| Immunogen | Mucins isolated from ovarian cyst fluid |
| Product Description | Recognizes a carbohydrate determinant of Gal 1-3(Fuc 1-4) GlcNAc which is blood group antigen Lewis A. It is present primarily on epithelial cells such as colon and kidneys. In the tumors and dedifferentiated tissues, decrease of Lewis A antigen was ... |
| Clonality | Monoclonal |
| Clone Designation | [7LE] |
| Molecular Weight | Multiple |
| Isotype | IgG1 |

| | |
|-------------------|---|
| NCBI | Not Applicable |
| Form | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Application Notes | Flow Cytometry (1-2ug/million cells), Immunofluorescence (1-2ug/ml), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min |