

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Thomsen-Friedenreich Antigen / CD176 (Pan Carcinoma Marker) Antibody, IgM, Clone: [A78-G/A7], Mouse, Monoclonal NBT-MSM1-936-P0

| | |
|----------------------------|---|
| Article Name | Thomsen-Friedenreich Antigen / CD176 (Pan Carcinoma Marker) Antibody, IgM, Clone: [A78-G/A7], Mouse, Monoclonal |
| Biozol Catalog Number | NBT-MSM1-936-P0 |
| Supplier Catalog Number | MSM1-936-P0 |
| Alternative Catalog Number | NBT-MSM1-936-P0-20,NBT-MSM1-936-P0-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Mouse |
| Category | Antikörper |
| Application | IF |
| Species Reactivity | Human, Mouse, Rat |
| Immunogen | Neuraminidase-treated human red blood cells |
| Product Description | Recognizes a disaccharide epitope, Gal 1-3GalNAc, of Thomsen-Friedenreich (TF) antigen. It is specific for both anomeric forms of the disaccharide (TF and TF , including related structures on the glycolipid) and shows no cross-reactivity with sialy... |
| Clonality | Monoclonal |
| Clone Designation | [A78-G/A7] |
| Molecular Weight | Multiple |
| Isotype | IgM |

| | |
|-------------------|---|
| NCBI | Not Applicable |
| Form | 200ug/ml of Ab purified from Bioreactor Concentrate. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Application Notes | Immunofluorescence (1-2ug/ml), Immunohistology (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes), |