

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

CDw17 (Lactosylceramide or LacCer) Antibody, IgM, Clone: [HO18.3G-6.F5], Mouse, Monoclonal NBT-MSM2-1092-P1BX

| | |
|----------------------------|---|
| Article Name | CDw17 (Lactosylceramide or LacCer) Antibody, IgM, Clone: [HO18.3G-6.F5], Mouse, Monoclonal |
| Biozol Catalog Number | NBT-MSM2-1092-P1BX |
| Supplier Catalog Number | MSM2-1092-P1BX |
| Alternative Catalog Number | NBT-MSM2-1092-P1BX-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Mouse |
| Category | Antikörper |
| Application | FC, IF |
| Species Reactivity | Human |
| Immunogen | beta-2 Microglobulin associated proteins from a detergent lysate of human PBL |
| Product Description | CD17 is an intermediate glycosphingolipid from the metabolism of higher gangliosides that localizes to sphingolipid-sterol rafts. CD17 is detectable in monocytes, granulocytes, basophils, platelets, a subset of peripheral B cells (CD19+) and tonsil d... |
| Clonality | Monoclonal |
| Clone Designation | [HO18.3G-6.F5] |
| Molecular Weight | Not Known |
| Isotype | IgM |

| | |
|-------------------|---|
| NCBI | Not known |
| Form | 200ug/ml of Ab Purified from Bioreactor Concentrate. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Application Notes | Flow Cytometry (1-2ug/million cells), Immunofluorescence (1-2ug/ml), Optimal dilution for a specific application should be determined. |