

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Cytokeratin 8/18 Antibody, IgG1, Clone: [KRT8.18/1346], Mouse, Monoclonal NBT-MSM27-1346-P0

| | |
|----------------------------|---|
| Article Name | Cytokeratin 8/18 Antibody, IgG1, Clone: [KRT8.18/1346], Mouse, Monoclonal |
| Biozol Catalog Number | NBT-MSM27-1346-P0 |
| Supplier Catalog Number | MSM27-1346-P0 |
| Alternative Catalog Number | NBT-MSM27-1346-P0-20,NBT-MSM27-1346-P0-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human |
| Immunogen | Cytoskeleton preparation from HeLa cells |
| Product Description | Cytokeratin 8 (CK8) belongs to the type II (or B or basic) subfamily of high molecular weight cytokeratins and exists in combination with cytokeratin 18 (CK18). This MAb recognizes all simple epithelia including glandular epithelium, for example thyr... |
| Clonality | Monoclonal |
| Clone Designation | [KRT8.18/1346] |
| Molecular Weight | 52.5kDa (CK8), 45kDa (CK18) |
| Isotype | IgG1 |

| | |
|-------------------|---|
| NCBI | 3856 |
| UniProt | P05787 |
| Form | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Application Notes | Immunohistochemistry (Formalin-fixed) (0.1-0.2ug/ml for 30 min at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes), Optimal dilution f |