

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Epstein-Barr Virus (LMP-1) Antibody, IgG1, Clone: [CS2], Mouse, Monoclonal NBT-MSM3-1742-P0

Article Name	Epstein-Barr Virus (LMP-1) Antibody, IgG1, Clone: [CS2], Mouse, Monoclonal
Biozol Catalog Number	NBT-MSM3-1742-P0
Supplier Catalog Number	MSM3-1742-P0
Alternative Catalog Number	NBT-MSM3-1742-P0-20,NBT-MSM3-1742-P0-100
Manufacturer	NeoBiotechnologies
Host	Mouse
Category	Antikörper
Application	ELISA, WB
Species Reactivity	Virus
Immunogen	Recombinant fusion protein containing the sequence of bacterial beta-galactosidase and the carboxyl half of EBV-encoded LMP
Product Description	This antibody is a mixture of four different monoclonal antibodies. This antibody is specific to 60kDa latent membrane protein (LMP-1) encoded by the BNLF1 gene of the EBV. Each clone reacts with different epitopes on the hydrophilic C-terminus of th
Clonality	Monoclonal
Clone Designation	[CS2]
Molecular Weight	60kDa
Isotype	lgG1

NCBI	Not Applicable
Form	200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prote
Antibody Type	Monoclonal Antibody
Application Notes	Immunofluorescence (0.5-1.0ug/ml), Western Blotting (0.5-1.0ug/ml),,Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min