

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Cytokeratin, pan (Epithelial Marker) Antibody, IgG2a, Clone: [PCK/3150], Mouse, Monoclonal NBT-MSM39-3150-P0

Article Name	Cytokeratin, pan (Epithelial Marker) Antibody, IgG2a, Clone: [PCK/3150], Mouse, Monoclonal
Biozol Catalog Number	NBT-MSM39-3150-P0
Supplier Catalog Number	MSM39-3150-P0
Alternative Catalog Number	NBT-MSM39-3150-P0-20,NBT-MSM39-3150-P0-100
Manufacturer	NeoBiotechnologies
Host	Mouse
Category	Antikörper
Application	FC, IF, IHC
Species Reactivity	Canine, Feline, Human
Immunogen	Crude cytokeratin extract prepared from RT-4 and MCF-7 cells.
Product Description	Anti-cytokeratin clone PCK/3150 demonstrates a broad spectrum of cytokeratin reactivity. In normal tissues, PCK/3150 is reactive with most epithelial types, including bile ducts and hepatocytes in liver, bladder epithelium, breast ducts, bronchial ep
Clonality	Monoclonal
Clone Designation	[PCK/3150]
Molecular Weight	40-67kDa (Multiple)
Isotype	IgG2a

NCBI	Multiple
Form	200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double 2005 azide. Also available WITHOUT BSA & Double 2005 azide at 1.0mg/ml.
Antibody Type	Monoclonal Antibody
Application Notes	Flow Cytometry (1-2ug/million cells), Immunofluorescence (1-2ug/ml),,Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 min at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at