

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Recombinant HSV1 (Herpes Simplex Virus Type I) Antibody, IgG1, Clone: [HSV1/1934], Mouse, Monoclonal NBT-MSM4-1934-P0

| | |
|----------------------------|---|
| Article Name | Recombinant HSV1 (Herpes Simplex Virus Type I) Antibody, IgG1, Clone: [HSV1/1934], Mouse, Monoclonal |
| Biozol Catalog Number | NBT-MSM4-1934-P0 |
| Supplier Catalog Number | MSM4-1934-P0 |
| Alternative Catalog Number | NBT-MSM4-1934-P0-20,NBT-MSM4-1934-P0-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Virus |
| Immunogen | Detergent-solubilized herpes simplex virus (HSV) type 1 infected cells |
| Product Description | The antibody reacts with HSV type 1 specific antigen. It is suitable for detection of HSV in human cellular material obtained from superficial lesions or biopsies and for the early identification of HSV in infected tissue cultures. The herpes simplex... |
| Clonality | Monoclonal |
| Clone Designation | [HSV1/1934] |
| Isotype | IgG1 |
| NCBI | Not Applicable |

| | |
|-------------------|---|
| Form | 200ug/ml of Ab Purified from rabbit anti-serum by Protein A. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA at 1.0mg/ml. |
| Antibody Type | Recombinant Monoclonal Antibody |
| Application Notes | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes), Optimal dilution f |