

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Recombinant Cytokeratin, Type I (Epithelial Marker) Antibody, IgG1, Clone: [rKRTL/6616], Mouse, Monoclonal NBT-MSM44-6616-P0

| | |
|----------------------------|---|
| Article Name | Recombinant Cytokeratin, Type I (Epithelial Marker) Antibody, IgG1, Clone: [rKRTL/6616], Mouse, Monoclonal |
| Biozol Catalog Number | NBT-MSM44-6616-P0 |
| Supplier Catalog Number | MSM44-6616-P0 |
| Alternative Catalog Number | NBT-MSM44-6616-P0-20,NBT-MSM44-6616-P0-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human |
| Immunogen | Synthetic peptide corresponding to Type I Cytokeratins residues within aa100-200 of Type I Cytokeratins |
| Product Description | The keratins are the typical intermediate filament proteins of epithelia, showing an outstanding degree of molecular diversity. Heteropolymeric filaments are formed by pairing of type I and type II molecules. In humans 54 functional keratin genes exi... |
| Clonality | Monoclonal |
| Clone Designation | [rKRTL/6616] |
| Molecular Weight | 60/53/51/45/40kDa |
| Isotype | IgG1 |

| | |
|-------------------|--|
| NCBI | 374454 |
| UniProt | Q7Z794 |
| Form | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Recombinant Monoclonal Antibody |
| Application Notes | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 20 minutes),Optimal diluti |