

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Melanoma Marker (MART-1 + Tyrosinase + gp100) Antibody, Clone: [DT101+ BC199 + T311 + HMB45], Mouse, Monoclonal NBT-MSM4-704-P0

| Article Name | Melanoma Marker (MART-1 + Tyrosinase + gp100) Antibody, Clone: [DT101+ BC199 + T311 + HMB45], Mouse, Monoclonal |
|----------------------------|---|
| Biozol Catalog Number | NBT-MSM4-704-P0 |
| Supplier Catalog Number | MSM4-704-P0 |
| Alternative Catalog Number | NBT-MSM4-704-P0-20,NBT-MSM4-704-P0-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human |
| Immunogen | Recombinant hMART-1 protein (DT101 & BC199), Extract of pigmented melanoma metastases from lymph nodes (HMB45), Recombinant tyrosinase protein (T311) |
| Product Description | This antibody cocktail recognizes three melanoma-specific proteins, which include MART-1, Tyrosinase and gp100. MART-1 is a newly identified melanocyte differentiation antigen recognized by autologous cytotoxic T lymphocytes. Tyrosinase is one of the |
| Clonality | Monoclonal |
| Clone Designation | [DT101+ BC199 + T311 + HMB45] |
| Molecular Weight | 20-22kDa (doublet) (MART-1): 90-100kDa (gp100), 70-80kDa (Tyrosinase) |

| NCBI | 2315 |
|-------------------|---|
| Form | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & amp; 0.05% azide. Also available WITHOUT BSA & amp; azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Application Notes | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes), Optimal dilution f |