

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Melanoma Marker (MART-1 + Tyrosinase + gp100) Antibody, Clone: [DT101+ BC199 + T311 + HMB45], Mouse, Monoclonal NBT-MSM4-704-P1ABX

Article Name	Melanoma Marker (MART-1 + Tyrosinase + gp100) Antibody, Clone: [DT101+ BC199 + T311 + HMB45], Mouse, Monoclonal
Biozol Catalog Number	NBT-MSM4-704-P1ABX
Supplier Catalog Number	MSM4-704-P1ABX
Alternative Catalog Number	NBT-MSM4-704-P1ABX-100
Manufacturer	NeoBiotechnologies
Host	Mouse
Category	Antikörper
Application	IHC
Species Reactivity	Human
Immunogen	Recombinant hMART-1 protein (DT101 & BC199), Extract of pigmented melanoma metastases from lymph nodes (HMB45), Recombinant tyrosinase protein (T311)
Product Description	This antibody cocktail recognizes three melanoma-specific proteins, which include MART-1, Tyrosinase and gp100. MART-1 is a newly identified melanocyte differentiation antigen recognized by autologous cytotoxic T lymphocytes. Tyrosinase is one of the
Clonality	Monoclonal
Clone Designation	[DT101+ BC199 + T311 + HMB45]
Molecular Weight	20-22kDa (doublet) (MART-1): 90-100kDa (gp100), 70-80kDa (Tyrosinase)

NCBI	2315
Form	200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Protein
Antibody Type	Monoclonal Antibody
Application Notes	Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes), Optimal dilution f