

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **HepPar-1 (Hepatocellular Marker) Antibody, IgG1, Clone: [HepPar1], Mouse, Monoclonal NBT-MSM4-966-P0**

Article Name	HepPar-1 (Hepatocellular Marker) Antibody, IgG1, Clone: [HepPar1], Mouse, Monoclonal
Biozol Catalog Number	NBT-MSM4-966-P0
Supplier Catalog Number	MSM4-966-P0
Alternative Catalog Number	NBT-MSM4-966-P0-20,NBT-MSM4-966-P0-100
Manufacturer	NeoBiotechnologies
Host	Mouse
Category	Antikörper
Application	IF, IHC
Species Reactivity	Canine, Human
Immunogen	Extract of a formalin-fixed, rejected-allograft of a human liver
Product Description	Hepatocyte Paraffin 1 or HepPar1 localizes to the mitochondria of hepatocytes. It is a sensitive marker for distinguishing hepatocellular carcinomas (HCC) from other metastatic carcinomas as well as cholangio-carcinomas. HCCs occur primarily in the s...
Clonality	Monoclonal
Clone Designation	[HepPar1]
Molecular Weight	Not Known
Isotype	IgG1

NCBI	Not Known
Form	200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Antibody Type	Monoclonal Antibody
Application Notes	Immunofluorescence (1-2ug/ml), Immunohistochemistry (Formalin-fixed) (0.25-0.5ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 1mM EDTA, pH 7.5-8.5, for 45 min at 95degC followed by cooling at RT for 20