

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Cytokeratin, Acidic (Type I or LMW) (Epithelial Marker) Antibody, IgG1, Clone: [SPM115], Mouse, Monoclonal NBT-MSM4X-253-P0

| Article Name               | Cytokeratin, Acidic (Type I or LMW) (Epithelial Marker) Antibody, IgG1, Clone: [SPM115], Mouse, Monoclonal   |
|----------------------------|--|
| Biozol Catalog Number      | NBT-MSM4X-253-P0   |
| Supplier Catalog Number    | MSM4X-253-P0   |
| Alternative Catalog Number | NBT-MSM4X-253-P0-20,NBT-MSM4X-253-P0-100   |
| Manufacturer               | NeoBiotechnologies   |
| Host                       | Mouse  |
| Category                   | Antikörper   |
| Application                | FC, IF, IHC, WB  |
| Species Reactivity         | Bovine, Canine, Gallus, Human, Monkey, Mouse, Rabbit, Rat, Reptile   |
| Immunogen                  | Human epidermal keratin  |
| Product Description        | This MAb recognizes the 56.5kDa (CK10), 50kDa (CK14), 50kDa (CK15), 48kDa (CK16), 40kDa (CK19) keratins of the acidic (Type I or LMW) subfamily. Twenty human keratins are resolved with two-dimensional gel electrophoresis into acidic (pl 6.0) subfamil |
| Clonality                  | Monoclonal   |
| Clone Designation          | [SPM115]   |
| Molecular Weight           | 56.5kDa (CK10), 50kDa (CK14), 50kDa (CK15), 48kDa (CK16), 40kDa (CK19)   |
| Isotype                    | IgG1   |

| NCBI              | 374454  |
|-------------------|---|
| UniProt           | Q7Z794  |
| Form              | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are a side. Also available WITHOUT BSA & Double at 1.0mg/ml.   |
| Antibody Type     | Monoclonal Antibody   |
| Application Notes | Flow Cytometry (1-2ug/million cells), Immunofluorescence (1-2ug/ml), Western Blot (1-2ug/ml), Immunohistochemistry (Formalinfixed) (1-2ug/ml for 30 min at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA |