

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Recombinant Cytokeratin, pan (Epithelial Marker) Antibody, IgG1, Clone: [Cocktail rPCK/6750], Mouse, Monoclonal NBT-MSM50-6750-P0

Article Name	Recombinant Cytokeratin, pan (Epithelial Marker) Antibody, IgG1, Clone: [Cocktail rPCK/6750], Mouse, Monoclonal
Biozol Catalog Number	NBT-MSM50-6750-P0
Supplier Catalog Number	MSM50-6750-P0
Alternative Catalog Number	NBT-MSM50-6750-P0-20,NBT-MSM50-6750-P0-100
Manufacturer	NeoBiotechnologies
Host	Mouse
Category	Antikörper
Application	FC, IF, IHC
Species Reactivity	Canine, Feline, Human
Immunogen	Crude cytokeratin extract prepared from RT-4 and MCF-7 cells.
Product Description	Anti-cytokeratin clone rPCK/6750 (CK rPCK/6750) demonstrates a broad spectrum of cytokeratin reactivity. In normal tissues, rPCK/6750 is reactive with most epithelial types, including bile ducts and hepatocytes in liver, bladder epithelium, breast du
Clonality	Monoclonal
Clone Designation	[Cocktail rPCK/6750]
Molecular Weight	40-67kDa (Multiple)
Isotype	IgG1

NCBI	Multiple
Form	200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are a side. Also available WITHOUT BSA & Double at 1.0mg/ml.
Antibody Type	Recombinant Monoclonal Antibody
Application Notes	Flow Cytometry (1-2ug/million cells), Immunofluorescence (1-2ug/ml),,Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min