

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Recombinant Hexa-histidine (Epitope Tag) Antibody, Clone: [6HIS/6402R], Rabbit, Monoclonal NBT-RBM2-6402-P0

| Article Name               | Recombinant Hexa-histidine (Epitope Tag) Antibody, Clone: [6HIS/6402R], Rabbit, Monoclonal   |
|----------------------------|--|
| Biozol Catalog Number      | NBT-RBM2-6402-P0   |
| Supplier Catalog Number    | RBM2-6402-P0   |
| Alternative Catalog Number | NBT-RBM2-6402-P0-20,NBT-RBM2-6402-P0-100   |
| Manufacturer               | NeoBiotechnologies   |
| Host                       | Rabbit   |
| Category                   | Antikörper   |
| Application                | FC, IF, WB   |
| Species Reactivity         | AII  |
| Immunogen                  | Hexa-histidine tagged human CD70 recombinant protein   |
| Product Description        | It recognizes both the free and protein-conjugated (either soluble or cell bound) form of hexa-histidine. This MAb is highly specific to hexa-histidine and shows no cross-reaction with other structurally related compounds. It has a very high affinity |
| Clonality                  | Monoclonal   |
| Clone Designation          | [6HIS/6402R]   |
| Molecular Weight           | 244Da  |
| NCBI                       | Not Applicable   |

| Form              | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prote |
|-------------------|--|
| Antibody Type     | Recombinant Monoclonal Antibody  |
| Application Notes | Confocal Microscopy, Western Blot (1-2ug/ml),,Flow Cytometry (0.5-1ug/million cells), Immunofluorescence (0.5-1ug/ml),, Immunoprecipitation (2-4ug/500ug proteins),,Optimal dilution for a specific application should be determined.  |