

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Recombinant Golgi Complex (Marker for Human Cells) Antibody, Clone: [GLG1/2829R], Rabbit, Monoclonal NBT-RBM3-2829-P1ABX

|                            |   |
|----------------------------|---|
| Article Name               | Recombinant Golgi Complex (Marker for Human Cells) Antibody, Clone: [GLG1/2829R], Rabbit, Monoclonal  |
| Biozol Catalog Number      | NBT-RBM3-2829-P1ABX   |
| Supplier Catalog Number    | RBM3-2829-P1ABX   |
| Alternative Catalog Number | NBT-RBM3-2829-P1ABX-100   |
| Manufacturer               | NeoBiotechnologies  |
| Host                       | Rabbit  |
| Category                   | Antikörper  |
| Application                | IHC   |
| Species Reactivity         | Human   |
| Immunogen                  | Recombinant full-length human GLG1 protein  |
| Product Description        | This MAb recognizes an antigen associated with the Golgi complex in human cells only. It can be used to stain the Golgi complex in cell or tissue preparations and can be used as a Golgi marker in subcellular fractions. It produces a diffuse staining ... |
| Clonality                  | Monoclonal  |
| Clone Designation          | [GLG1/2829R]  |
| Molecular Weight           | 134kDa  |
| NCBI                       | <a href="#">2734</a>  |

|                   |   |
|-------------------|---|
| UniProt           | <a href="#">Q92896</a>  |
| Form              | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA at 1.0mg/ml.  |
| Antibody Type     | Recombinant Monoclonal Antibody   |
| Application Notes | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes), Optimal dilution f |