

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Mouse IgG F(c) Antibody Peroxidase Conjugated, HRP, Goat, Polyclonal ROC-610-1303

| Article Name Mouse IgG F(c) Antibody Peroxidase Conjugated, HRP, Goat, Polyclonal Biozol Catalog Number ROC-610-1303 Supplier Catalog Number 610-1303 Alternative Catalog Number ROC-610-1303 Manufacturer Rockland Immunochemicals Host Goat Category Antikörper Application ELISA, WB Species Reactivity Mouse Immunogen Anti-Mouse IgG F(c) fragment was produced by repeated immunization with Mouse IgG F(c) fragment in goat. Conjugation HRP Clonality Polyclonal Concentration 2.0 mg/mL by UV absorbance at 280 nm Buffer 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 Form Lyophilized Target Mouse Antibody Type Secondary Antibody | | |
|--|----------------------------|--|
| Supplier Catalog Number 610-1303 Alternative Catalog Number ROC-610-1303 Manufacturer Rockland Immunochemicals Host Goat Category Antikörper Application ELISA, WB Species Reactivity Mouse Immunogen Anti-Mouse IgG F(c) fragment was produced by repeated immunization with Mouse IgG F(c) fragment in goat. Conjugation HRP Clonality Polyclonal Concentration 2.0 mg/mL by UV absorbance at 280 nm Buffer 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 Form Lyophilized Target Mouse | Article Name | |
| Alternative Catalog Number ROC-610-1303 Manufacturer Rockland Immunochemicals Host Goat Category Antikörper Application ELISA, WB Species Reactivity Mouse Immunogen Anti-Mouse IgG F(c) fragment was produced by repeated immunization with Mouse IgG F(c) fragment in goat. Conjugation HRP Clonality Polyclonal Concentration 2.0 mg/mL by UV absorbance at 280 nm Buffer 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 Form Lyophilized Target Mouse | Biozol Catalog Number | ROC-610-1303 |
| Manufacturer Rockland Immunochemicals Host Goat Category Antikörper Application ELISA, WB Species Reactivity Mouse Immunogen Anti-Mouse IgG F(c) fragment was produced by repeated immunization with Mouse IgG F(c) fragment in goat. Conjugation HRP Clonality Polyclonal Concentration 2.0 mg/mL by UV absorbance at 280 nm Buffer 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 Form Lyophilized Target Mouse | Supplier Catalog Number | 610-1303 |
| Host Goat Category Antikörper Application ELISA, WB Species Reactivity Mouse Immunogen Anti-Mouse IgG F(c) fragment was produced by repeated immunization with Mouse IgG F(c) fragment in goat. Conjugation HRP Clonality Polyclonal Concentration 2.0 mg/mL by UV absorbance at 280 nm Buffer 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 Form Lyophilized Target Mouse | Alternative Catalog Number | ROC-610-1303 |
| Category Antikörper Application ELISA, WB Species Reactivity Mouse Immunogen Anti-Mouse IgG F(c) fragment was produced by repeated immunization with Mouse IgG F(c) fragment in goat. Conjugation HRP Clonality Polyclonal Concentration 2.0 mg/mL by UV absorbance at 280 nm Buffer 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 Form Lyophilized Target Mouse | Manufacturer | Rockland Immunochemicals |
| Application ELISA, WB Species Reactivity Mouse Immunogen Anti-Mouse IgG F(c) fragment was produced by repeated immunization with Mouse IgG F(c) fragment in goat. Conjugation HRP Clonality Polyclonal Concentration 2.0 mg/mL by UV absorbance at 280 nm Buffer 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 Form Lyophilized Target Mouse | Host | Goat |
| Species Reactivity Immunogen Anti-Mouse IgG F(c) fragment was produced by repeated immunization with Mouse IgG F(c) fragment in goat. Conjugation HRP Clonality Polyclonal Concentration 2.0 mg/mL by UV absorbance at 280 nm Buffer 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 Form Lyophilized Target Mouse | Category | Antikörper |
| Immunogen Anti-Mouse IgG F(c) fragment was produced by repeated immunization with Mouse IgG F(c) fragment in goat. Conjugation HRP Clonality Polyclonal Concentration 2.0 mg/mL by UV absorbance at 280 nm Buffer 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 Form Lyophilized Target Mouse | Application | ELISA, WB |
| immunization with Mouse IgG F(c) fragment in goat. Conjugation HRP Clonality Polyclonal Concentration 2.0 mg/mL by UV absorbance at 280 nm Buffer 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 Form Lyophilized Target Mouse | Species Reactivity | Mouse |
| Clonality Polyclonal Concentration 2.0 mg/mL by UV absorbance at 280 nm Buffer 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 Form Lyophilized Target Mouse | Immunogen | |
| Concentration 2.0 mg/mL by UV absorbance at 280 nm Buffer 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 Form Lyophilized Target Mouse | Conjugation | HRP |
| Buffer 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 Form Lyophilized Target Mouse | Clonality | Polyclonal |
| Form Lyophilized Target Mouse | Concentration | 2.0 mg/mL by UV absorbance at 280 nm |
| Target Mouse | Buffer | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
| | Form | Lyophilized |
| Antibody Type Secondary Antibody | Target | Mouse |
| | Antibody Type | Secondary Antibody |

| Application Dilute | ELISA: 1:10,000 - 1:50,000, IHC: 1:500 - 1:2,500, WB: 1:1,000 - 1:5,000 |
|--------------------|---|
| Application Notes | Anti-Mouse IgG F(c) fragment Peroxidase conjugated Antibody has been tested by ELISA and western blot and is suitable for use in immunoelectrophoresis, western-blot, competitive western-blot, ELISA and competitive ELISA assays. Specific conditions for re |