

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | Fax: +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-Hu CD319 PE, Clone: [162], Monoclonal EXB-1P-908-T100

Article Name Anti-Hu CD319 PE, Clone: [162], Monoclonal Biozol Catalog Number EXB-1P-908-T100 Supplier Catalog Number 1P-908-T100 Alternative Catalog Number EXB-1P-908-T100 Manufacturer EXBIO Category Antikörper Application FC Species Reactivity Human Immunogen Recombinant extracellular domain of human CD319 (204 aa) fused with Fc part of human IgG1 Conjugation PE CD319 is a homophilic type I transmembrane protein with two intracellular immunoreceptor tyrosine-based switch motives (ITSMs). It is expressed by cytotoxic T cells, activated B cells, NK cells, and mature dendritic cells. CD319 triggers NK cell-medi Clonality Monoclonal Clone Designation [162] Buffer Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide Target CD319		
Supplier Catalog Number 1P-908-T100 Alternative Catalog Number EXB-1P-908-T100 Manufacturer EXBIO Category Antikörper Application FC Species Reactivity Human Immunogen Recombinant extracellular domain of human CD319 (204 aa) fused with Fc part of human IgG1 Conjugation PE CD319 is a homophilic type I transmembrane protein with two intracellular immunoreceptor tyrosine-based switch motives (ITSMs). It is expressed by cytotoxic T cells, activated B cells, NK cells, and mature dendritic cells. CD319 triggers NK cell-medi Clonality Monoclonal Clone Designation [162] Buffer Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Article Name	Anti-Hu CD319 PE, Clone: [162], Monoclonal
Alternative Catalog Number EXB-1P-908-T100 Manufacturer EXBIO Category Antikörper Application FC Species Reactivity Human Immunogen Recombinant extracellular domain of human CD319 (204 aa) fused with Fc part of human IgG1 Conjugation PE CD319 is a homophilic type I transmembrane protein with two intracellular immunoreceptor tyrosine-based switch motives (ITSMs). It is expressed by cytotoxic T cells, activated B cells, NK cells, and mature dendritic cells. CD319 triggers NK cell-medi Clone Designation [162] Buffer Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Biozol Catalog Number	EXB-1P-908-T100
Manufacturer EXBIO Category Antikörper Application FC Species Reactivity Human Recombinant extracellular domain of human CD319 (204 aa) fused with Fc part of human IgG1 Conjugation PE CD319 is a homophilic type I transmembrane protein with two intracellular immunoreceptor tyrosine-based switch motives (ITSMs). It is expressed by cytotoxic T cells, activated B cells, NK cells, and mature dendritic cells. CD319 triggers NK cell-medi Clonality Monoclonal Clone Designation [162] Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Supplier Catalog Number	1P-908-T100
Category Antikörper Application FC Species Reactivity Human Recombinant extracellular domain of human CD319 (204 aa) fused with Fc part of human IgG1 Conjugation PE CD319 is a homophilic type I transmembrane protein with two intracellular immunoreceptor tyrosine-based switch motives (ITSMs). It is expressed by cytotoxic T cells, activated B cells, NK cells, and mature dendritic cells. CD319 triggers NK cell-medi Clonality Monoclonal Clone Designation [162] Buffer Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Alternative Catalog Number	EXB-1P-908-T100
Application FC Species Reactivity Human Recombinant extracellular domain of human CD319 (204 aa) fused with Fc part of human IgG1 Conjugation PE CD319 is a homophilic type I transmembrane protein with two intracellular immunoreceptor tyrosine-based switch motives (ITSMs). It is expressed by cytotoxic T cells, activated B cells, NK cells, and mature dendritic cells. CD319 triggers NK cell-medi Clonality Monoclonal Clone Designation [162] Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Manufacturer	EXBIO
Species Reactivity Human Recombinant extracellular domain of human CD319 (204 aa) fused with Fc part of human IgG1 Conjugation PE CD319 is a homophilic type I transmembrane protein with two intracellular immunoreceptor tyrosine-based switch motives (ITSMs). It is expressed by cytotoxic T cells, activated B cells, NK cells, and mature dendritic cells. CD319 triggers NK cell-medi Clonality Monoclonal Clone Designation [162] Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Category	Antikörper
Immunogen Recombinant extracellular domain of human CD319 (204 aa) fused with Fc part of human IgG1 Conjugation PE CD319 is a homophilic type I transmembrane protein with two intracellular immunoreceptor tyrosine-based switch motives (ITSMs). It is expressed by cytotoxic T cells, activated B cells, NK cells, and mature dendritic cells. CD319 triggers NK cell-medi Clonality Monoclonal Clone Designation [162] Buffer Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Application	FC
mmunogen with Fc part of human IgG1 Conjugation PE CD319 is a homophilic type I transmembrane protein with two intracellular immunoreceptor tyrosine-based switch motives (ITSMs). It is expressed by cytotoxic T cells, activated B cells, NK cells, and mature dendritic cells. CD319 triggers NK cell-medi Clonality Monoclonal Clone Designation [162] Buffer Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Species Reactivity	Human
Product Description CD319 is a homophilic type I transmembrane protein with two intracellular immunoreceptor tyrosine-based switch motives (ITSMs). It is expressed by cytotoxic T cells, activated B cells, NK cells, and mature dendritic cells. CD319 triggers NK cell-medi Clonality Monoclonal Clone Designation [162] Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Immunogen	,
Product Description intracellular immunoreceptor tyrosine-based switch motives (ITSMs). It is expressed by cytotoxic T cells, activated B cells, NK cells, and mature dendritic cells. CD319 triggers NK cell-medi Clonality Monoclonal Clone Designation [162] Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Conjugation	PE
Clone Designation [162] Buffer Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Product Description	intracellular immunoreceptor tyrosine-based switch motives (ITSMs). It is expressed by cytotoxic T cells, activated B cells, NK cells, and
Buffer Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Clonality	Monoclonal
azide	Clone Designation	[162]
Target CD319	Buffer	
	Target	CD319

۸.				NIatoo
М	μμ	IICali	OH	Notes

Flow cytometry: The reagent is designed for analysis of human blood cells using 10 μ l reagent / 100 μ l of whole blood or 106 cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.