

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Anti-mTOR/MTOR Antibody Picoband(TM), Rabbit, Polyclonal BOB-A00003-2

|                            |  |
|----------------------------|--|
| Article Name               | Anti-mTOR/MTOR Antibody Picoband(TM), Rabbit, Polyclonal   |
| Biozol Catalog Number      | BOB-A00003-2   |
| Supplier Catalog Number    | A00003-2   |
| Alternative Catalog Number | BOB-A00003-2-10UG,BOB-A00003-2-100UG   |
| Manufacturer               | Boster Bio   |
| Host                       | Rabbit   |
| Category                   | Antikörper   |
| Application                | ELISA, IHC, WB   |
| Species Reactivity         | Human, Monkey  |
| Immunogen                  | E.coli-derived human mTOR/MTOR recombinant protein (Position: N2093-N2537).  |
| Product Description        | Boster Bio Anti-mTOR/MTOR Antibody Picoband(TM) catalog A00003-2. Tested in ELISA, IHC, WB applications. This antibody reacts with Human, Monkey.... |
| Clonality                  | Polyclonal   |
| Concentration              | Adding 0.2 ml of distilled water will yield a concentration of 500 µg/ml.  |
| Molecular Weight           | 289 kDa  |
| UniProt                    | <a href="#">P42345</a>   |

|                    |  |
|--------------------|--|
| Buffer             | Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na <sub>2</sub> HPO <sub>4</sub> , 0.05mg NaN <sub>3</sub> .   |
| Purity             | Immunogen affinity purified.   |
| Form               | Lyophilized  |
| Application Dilute | Western blot, 0.1-0.5µg/ml Immunohistochemistry (Paraffin-embedded Section), 0.5-1µg/ml Direct ELISA, 0.1-0.5µg/ml   |
| Application Notes  | Tested Species: In-house tested species with positive results. By Heat: Boiling the paraffin sections in 10mM citrate buffer, pH6.0, for 20mins is required for the staining of formalin/paraffin sections. Other applications have not been tested. Optimal d |