

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-CD3 epsilon/CD3E Antibody Picoband(TM), Rabbit, Polyclonal BOB-PB9093

Article Name	Anti-CD3 epsilon/CD3E Antibody Picoband(TM), Rabbit, Polyclonal
Biozol Catalog Number	BOB-PB9093
Supplier Catalog Number	PB9093
Alternative Catalog Number	BOB-PB9093-10UG,BOB-PB9093-100UG
Manufacturer	Boster Bio
Host	Rabbit
Category	Antikörper
Application	ICC, IF, IHC, IHC-Fr, WB
Species Reactivity	Gallus, Human, Mouse, Rat
Immunogen	E.coli-derived human CD3 epsilon recombinant protein (Position: D23-I207). Human CD3 epsilon shares 65% amino acid (aa) sequence identity with mouse CD3 epsilon.
Product Description	Boster Bio Anti-CD3 epsilon/CD3E Antibody Picoband(TM) catalog PB9093. Tested in IF, IHC, IHC-F, ICC, WB applications. This antibody reacts with Chicken, Human, Mouse, Rat
Clonality	Polyclonal
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 $\mu g/ml$.
Molecular Weight	23147 MW
UniProt	P07766

Buffer	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.
Purity	Immunogen affinity purified.
Form	Lyophilized
Application Dilute	Western blot, 0.1 - $0.5\mu g/ml$, Human Immunohistochemistry (Paraffinembedded Section), 0.5 - $1\mu g/ml$, Human, Mouse, Rat, Chicken, By Heat Immunohistochemistry (Frozen Section), 0.5 - $1\mu g/ml$, Mouse, Rat Immunocytochemistry , 0.5 - $1\mu g/ml$, Human Immu
Application Notes	WB: The detection limit for CD3 epsilon is approximately 0.25ng/lane under reducing conditions. Tested Species: In-house tested species with positive results. By Heat: Boiling the paraffin sections in 10mM citrate buffer, pH6.0, for 20mins is required fo