

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-Interleukin-10 IL10 Antibody, Rabbit, Polyclonal BOB-RP1014

| | |
|----------------------------|---|
| Article Name | Anti-Interleukin-10 IL10 Antibody, Rabbit, Polyclonal |
| Biozol Catalog Number | BOB-RP1014 |
| Supplier Catalog Number | RP1014 |
| Alternative Catalog Number | BOB-RP1014-10UG,BOB-RP1014-100UG |
| Manufacturer | Boster Bio |
| Host | Rabbit |
| Category | Antikörper |
| Application | ELISA, IHC, WB |
| Species Reactivity | Human |
| Immunogen | E. coli-derived human IL-10 recombinant protein (Position: S19-N178). |
| Product Description | Boster Bio Anti-Interleukin-10 IL10 Antibody catalog RP1014. Tested in ELISA, IHC, WB applications. This antibody reacts with Human.... |
| Clonality | Polyclonal |
| Concentration | Adding 0.2 ml of distilled water will yield a concentration of 500 µg/ml. |
| Molecular Weight | 20517 MW |
| UniProt | P22301 |

| | |
|--------------------|---|
| Buffer | Each vial contains 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ . Carrier free (No BSA) form available in stock. If you want this antibody carrier free please specify Carrier Free or No BSA in your order note. |
| Purity | Immunogen affinity purified. |
| Form | Lyophilized |
| Application Dilute | Immunohistochemistry (Paraffin-embedded Section), 0.5-1µg/ml, Human, By Heat ELISA , 0.1-0.5µg/ml, Human, -Western blot, 0.1-0.5µg/ml, Human |
| Application Notes | By Heat: Boiling the paraffin sections in 10mM citrate buffer, pH6.0, for 20mins is required for the staining of formalin/paraffin sections. Other applications have not been tested. Optimal dilutions should be determined by end users. |