

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

HDAC2 Antibody, Unconjugated, Rabbit, Polyclonal PRS-7899

| Article Name | HDAC2 Antibody, Unconjugated, Rabbit, Polyclonal |
|----------------------------|--|
| Biozol Catalog Number | PRS-7899 |
| Supplier Catalog Number | 7899 |
| Alternative Catalog Number | PRS-7899-0.02, PRS-7899-0.1 |
| Manufacturer | ProSci |
| Host | Rabbit |
| Category | Antikörper |
| Application | ELISA, IF, IHC-P, WB |
| Species Reactivity | Human, Mouse, Rat |
| Immunogen | HDAC2 antibody was raised against an 18 amino acid peptide near the carboxy terminus of human HDAC2. The immunogen is located within the last 50 amino acids of HDAC2. |
| Conjugation | Unconjugated |
| Clonality | Polyclonal |
| Concentration | 1 mg/mL |
| Molecular Weight | Predicted: 54 kDa Observed: 52 kDa |
| NCBI | 3066 |
| UniProt | Q92769 |
| Buffer | HDAC2 antibody is supplied in PBS containing 0.02% sodium azide. |

| Form | Liquid |
|--------------------|---|
| Application Dilute | Optimal dilutions for each application to be determined by the researcher. |
| Application Notes | HDAC2 antibody can be used for detection of HDAC2 by Western blot at 0.5 - 1 μ g/ml. Antibody can also be used for Immunohistochemistry at 5 μ g/mL. For Immunoflorescence start at 20 μ g/mL.Antibody validated: Western Blot in human samples, Immunohistoc |