

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

EXTL2 Polyclonal Antibody, Biotin Conjugated, Rabbit BSS-BS-6005R-BIOTIN

Article Name EXTL2 Polyclonal Antibody, Biotin Conjugated, Rabbit Biozol Catalog Number BSS-BS-6005R-BIOTIN Alternative Catalog Number BSS-BS-6005R-BIOTIN Alternative Catalog Number BSS-BS-6005R-BIOTIN-100 Manufacturer Bioss Host Rabbit Category Antikörper Application ELISA, IHC-Fr, IHC-P Immunogen 5-110/330 Conjugation Biotin EXTL2 is a glycosyltransferase required for the biosynthesis of heparan-sulfate and is responsible for the alternating addition of beta-1-4-linked glucuronic acid (GIcA) and alpha-1-4-linked Nacetylglucosamine (GIcNAc) units to nascent heparan sulfa Clonality Polyclonal Concentration 1ug/ul NCBI 2134 Buffer Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Source KLH conjugated synthetic peptide derived from human EXTL		
Supplier Catalog Number BS-6005R-BIOTIN Alternative Catalog Number BSS-BS-6005R-BIOTIN-100 Manufacturer Bioss Host Rabbit Category Antikörper Application ELISA, IHC-Fr, IHC-P Immunogen 5-110/330 Conjugation Biotin Product Description EXTL2 is a glycosyltransferase required for the biosynthesis of heparan-sulfate and is responsible for the alternating addition of beta-1-4-linked glucuronic acid (GlcA) and alpha-1-4-linked N-acetylglucosamine (GlcNAc) units to nascent heparan sulfa Clonality Polyclonal Concentration 1ug/ul NCBI 2134 Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.	Article Name	EXTL2 Polyclonal Antibody, Biotin Conjugated, Rabbit
Alternative Catalog Number BSS-BS-6005R-BIOTIN-100 Manufacturer Bioss Host Rabbit Category Antikörper Application ELISA, IHC-Fr, IHC-P Immunogen 5-110/330 Conjugation Biotin EXTL2 is a glycosyltransferase required for the biosynthesis of heparan-sulfate and is responsible for the alternating addition of beta-1-4-linked glucuronic acid (GIcA) and alpha-1-4-linked Nacetylglucosamine (GIcNAc) units to nascent heparan sulfa Clonality Polyclonal Concentration 1ug/ul NCBI Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.	Biozol Catalog Number	BSS-BS-6005R-BIOTIN
Manufacturer Bioss Host Rabbit Category Antikörper Application ELISA, IHC-Fr, IHC-P Immunogen 5-110/330 Conjugation Biotin EXTL2 is a glycosyltransferase required for the biosynthesis of heparan-sulfate and is responsible for the alternating addition of beta-1-4-linked glucuronic acid (GlcA) and alpha-1-4-linked Nacetylglucosamine (GlcNAc) units to nascent heparan sulfa Clonality Polyclonal Concentration 1ug/ul NCBI 2134 Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.	Supplier Catalog Number	BS-6005R-BIOTIN
Host Rabbit Category Antikörper Application ELISA, IHC-Fr, IHC-P Immunogen 5-110/330 Conjugation Biotin EXTL2 is a glycosyltransferase required for the biosynthesis of heparan-sulfate and is responsible for the alternating addition of beta-1-4-linked glucuronic acid (GlcA) and alpha-1-4-linked N-acetylglucosamine (GlcNAc) units to nascent heparan sulfa Clonality Polyclonal Concentration 1ug/ul NCBI 2134 Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.	Alternative Catalog Number	BSS-BS-6005R-BIOTIN-100
Category Antikörper Application ELISA, IHC-Fr, IHC-P Immunogen 5-110/330 Conjugation Biotin EXTL2 is a glycosyltransferase required for the biosynthesis of heparan-sulfate and is responsible for the alternating addition of beta-1-4-linked glucuronic acid (GlcA) and alpha-1-4-linked Nacetylglucosamine (GlcNAc) units to nascent heparan sulfa Clonality Polyclonal Concentration 1ug/ul NCBI 2134 Buffer Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.	Manufacturer	Bioss
Application ELISA, IHC-Fr, IHC-P Immunogen 5-110/330 Conjugation Biotin EXTL2 is a glycosyltransferase required for the biosynthesis of heparan-sulfate and is responsible for the alternating addition of beta-1-4-linked glucuronic acid (GlcA) and alpha-1-4-linked N-acetylglucosamine (GlcNAc) units to nascent heparan sulfa Clonality Polyclonal Concentration 1ug/ul NCBI 2134 Buffer Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.	Host	Rabbit
Immunogen 5-110/330 Conjugation Biotin EXTL2 is a glycosyltransferase required for the biosynthesis of heparan-sulfate and is responsible for the alternating addition of beta-1-4-linked glucuronic acid (GlcA) and alpha-1-4-linked Nacetylglucosamine (GlcNAc) units to nascent heparan sulfa Clonality Polyclonal Concentration lug/ul NCBI 2134 Buffer Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.	Category	Antikörper
Conjugation Biotin EXTL2 is a glycosyltransferase required for the biosynthesis of heparan-sulfate and is responsible for the alternating addition of beta-1-4-linked glucuronic acid (GlcA) and alpha-1-4-linked N-acetylglucosamine (GlcNAc) units to nascent heparan sulfa Clonality Polyclonal Concentration 1ug/ul NCBI 2134 Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.	Application	ELISA, IHC-Fr, IHC-P
Product Description EXTL2 is a glycosyltransferase required for the biosynthesis of heparan-sulfate and is responsible for the alternating addition of beta-1-4-linked glucuronic acid (GlcA) and alpha-1-4-linked N-acetylglucosamine (GlcNAc) units to nascent heparan sulfa Clonality Polyclonal Concentration 1ug/ul NCBI 2134 Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.	Immunogen	5-110/330
heparan-sulfate and is responsible for the alternating addition of beta-1-4-linked glucuronic acid (GlcA) and alpha-1-4-linked N-acetylglucosamine (GlcNAc) units to nascent heparan sulfa Clonality Polyclonal Concentration lug/ul NCBI 2134 Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.	Conjugation	Biotin
Concentration 1ug/ul NCBI 2134 Buffer Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.	Product Description	heparan-sulfate and is responsible for the alternating addition of beta-1-4-linked glucuronic acid (GlcA) and alpha-1-4-linked N-
NCBI 2134 Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.	Clonality	Polyclonal
Buffer Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.	Concentration	1ug/ul
BSA, 0.02% Proclin300 and 50% Glycerol.	NCBI	2134
Source KLH conjugated synthetic peptide derived from human EXTL	Buffer	•
	Source	KLH conjugated synthetic peptide derived from human EXTL

Purity	Purified by Protein A.
Target	EXTL2
Application Dilute	ELISA(1:500-1000), IHC-P(1:200-400), IHC-F(1:100-500)