

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

CD3g Antibody / CD3 gamma, Rabbit, Polyclonal NSJ-RQ8022

Article Name CD3g Antibody / CD3 gamma, Rabbit, Polyclonal Biozol Catalog Number NSJ-RQ8022 Alternative Catalog Number NSJ-RQ8022 Manufacturer NSJ Bioreagents Host Rabbit Category Antikörper Application ELISA, FACS, IF, WB Species Reactivity Human Immunogen E. coli-derived recombinant human protein (amino acids Q23-D165) was used as the immunogen for the CD3 gamma antibody. Product Description Product Description Product Description Polyclonal UniProt P09693 Purity Antigen affinity purified Form 0.5mg/ml if reconstituted with 0.2ml sterile DI water Antibody Type Primary Antibody		
Supplier Catalog Number RQ8022 Alternative Catalog Number NSJ-RQ8022 Manufacturer NSJ Bioreagents Host Rabbit Category Antikörper Application ELISA, FACS, IF, WB Species Reactivity Human Immunogen E. coli-derived recombinant human protein (amino acids Q23-D165) was used as the immunogen for the CD3 gamma antibody. Product Description Product Description Product Description Product Description Polyclonal UniProt Po9693 Purity Antigen affinity purified Form 0.5mg/ml if reconstituted with 0.2ml sterile DI water	Article Name	CD3g Antibody / CD3 gamma, Rabbit, Polyclonal
Alternative Catalog Number NSJ-RQ8022 Manufacturer NSJ Bioreagents Host Rabbit Category Antikörper Application ELISA, FACS, IF, WB Species Reactivity Human E. coli-derived recombinant human protein (amino acids Q23-D165) was used as the immunogen for the CD3 gamma antibody. T-cell surface glycoprotein CD3 gamma chain is a protein that in humans is encoded by the CD3G gene. The protein encoded by this gene is the CD3-gamma polypeptide, which together with CD3-epsilon, -delta and -zeta, and the T-cell receptor alpha/beta Clonality Polyclonal UniProt P09693 Purity Antigen affinity purified Form 0.5mg/ml if reconstituted with 0.2ml sterile DI water	Biozol Catalog Number	NSJ-RQ8022
Manufacturer NSJ Bioreagents Host Rabbit Category Antikörper Application ELISA, FACS, IF, WB Species Reactivity Human E. coli-derived recombinant human protein (amino acids Q23-D165) was used as the immunogen for the CD3 gamma antibody. T-cell surface glycoprotein CD3 gamma chain is a protein that in humans is encoded by the CD3G gene. The protein encoded by this gene is the CD3-gamma polypeptide, which together with CD3-epsilon, -delta and -zeta, and the T-cell receptor alpha/beta Clonality Polyclonal UniProt Po9693 Purity Antigen affinity purified Form 0.5mg/ml if reconstituted with 0.2ml sterile DI water	Supplier Catalog Number	RQ8022
Host Rabbit Category Antikörper Application ELISA, FACS, IF, WB Species Reactivity Human E. coli-derived recombinant human protein (amino acids Q23-D165) was used as the immunogen for the CD3 gamma antibody. T-cell surface glycoprotein CD3 gamma chain is a protein that in humans is encoded by the CD3G gene. The protein encoded by this gene is the CD3-gamma polypeptide, which together with CD3-epsilon, -delta and -zeta, and the T-cell receptor alpha/beta Clonality Polyclonal UniProt P09693 Purity Antigen affinity purified Form 0.5mg/ml if reconstituted with 0.2ml sterile DI water	Alternative Catalog Number	NSJ-RQ8022
Category Antikörper Application ELISA, FACS, IF, WB Species Reactivity Human E. coli-derived recombinant human protein (amino acids Q23-D165) was used as the immunogen for the CD3 gamma antibody. T-cell surface glycoprotein CD3 gamma chain is a protein that in humans is encoded by the CD3G gene. The protein encoded by this gene is the CD3-gamma polypeptide, which together with CD3-epsilon, -delta and -zeta, and the T-cell receptor alpha/beta Clonality Polyclonal UniProt P09693 Purity Antigen affinity purified Form 0.5mg/ml if reconstituted with 0.2ml sterile DI water	Manufacturer	NSJ Bioreagents
Application ELISA, FACS, IF, WB Species Reactivity Human E. coli-derived recombinant human protein (amino acids Q23-D165) was used as the immunogen for the CD3 gamma antibody. T-cell surface glycoprotein CD3 gamma chain is a protein that in humans is encoded by the CD3G gene. The protein encoded by this gene is the CD3-gamma polypeptide, which together with CD3-epsilon, -delta and -zeta, and the T-cell receptor alpha/beta Clonality Polyclonal UniProt P09693 Purity Antigen affinity purified Form 0.5mg/ml if reconstituted with 0.2ml sterile DI water	Host	Rabbit
Species Reactivity Human E. coli-derived recombinant human protein (amino acids Q23-D165) was used as the immunogen for the CD3 gamma antibody. T-cell surface glycoprotein CD3 gamma chain is a protein that in humans is encoded by the CD3G gene. The protein encoded by this gene is the CD3-gamma polypeptide, which together with CD3-epsilon, -delta and -zeta, and the T-cell receptor alpha/beta Clonality Polyclonal UniProt P09693 Purity Antigen affinity purified Form 0.5mg/ml if reconstituted with 0.2ml sterile DI water	Category	Antikörper
Immunogen E. coli-derived recombinant human protein (amino acids Q23-D165) was used as the immunogen for the CD3 gamma antibody. T-cell surface glycoprotein CD3 gamma chain is a protein that in humans is encoded by the CD3G gene. The protein encoded by this gene is the CD3-gamma polypeptide, which together with CD3-epsilon, -delta and -zeta, and the T-cell receptor alpha/beta Clonality Polyclonal UniProt Po9693 Purity Antigen affinity purified Form 0.5mg/ml if reconstituted with 0.2ml sterile DI water	Application	ELISA, FACS, IF, WB
was used as the immunogen for the CD3 gamma antibody. T-cell surface glycoprotein CD3 gamma chain is a protein that in humans is encoded by the CD3G gene. The protein encoded by this gene is the CD3-gamma polypeptide, which together with CD3-epsilon, -delta and -zeta, and the T-cell receptor alpha/beta Clonality Polyclonal UniProt P09693 Purity Antigen affinity purified Form 0.5mg/ml if reconstituted with 0.2ml sterile DI water	Species Reactivity	Human
Product Description humans is encoded by the CD3G gene. The protein encoded by this gene is the CD3-gamma polypeptide, which together with CD3-epsilon, -delta and -zeta, and the T-cell receptor alpha/beta Clonality Polyclonal UniProt P09693 Purity Antigen affinity purified Form 0.5mg/ml if reconstituted with 0.2ml sterile DI water	Immunogen	·
UniProt P09693 Purity Antigen affinity purified Form 0.5mg/ml if reconstituted with 0.2ml sterile DI water	Product Description	humans is encoded by the CD3G gene. The protein encoded by this gene is the CD3-gamma polypeptide, which together with CD3-
Purity Antigen affinity purified Form 0.5mg/ml if reconstituted with 0.2ml sterile DI water	Clonality	Polyclonal
Form 0.5mg/ml if reconstituted with 0.2ml sterile DI water	UniProt	P09693
	Purity	Antigen affinity purified
Antibody Type Primary Antibody	Form	0.5mg/ml if reconstituted with 0.2ml sterile DI water
	Antibody Type	Primary Antibody

Application Dilute

Western blot: 0.5-1ug/ml,Immunofluorescence: 5ug/ml,Flow

cytometry: 1-3ug/million cells,Direct ELISA: 0.1-0.5ug/ml