

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

NFKB2 Antibody, Rabbit, Polyclonal ASB-OASG04979

Article Name NFKB2 Antibody, Rabbit, Polyclonal Biozol Catalog Number ASB-OASG04979 Supplier Catalog Number OASG04979 Alternative Catalog Number ASB-OASG04979-100UL Manufacturer Aviva Host Rabbit Category Antikörper Application IF, IHC, IP, WB Species Reactivity Human, Mouse, Rat Immunogen Synthesized peptide derived from human NFKappaB-p100 around the non-phosphorylation site of S865. Location: 810-890aa. This gene encodes a subunit of the transcription factor complex nuclear factor-kappa-B (NFkB). The NFkB complex is expressed in numerous cell types and functions as a central activator of genes involved in inflammation and immune function. The protei Clonality Polyclonal Concentration 1 mg/ml Molecular Weight 97 kDa NCBI 4791 UniProt Q00653, Q9WTK5		
Supplier Catalog Number ASB-OASG04979 Alternative Catalog Number ASB-OASG04979-100UL Manufacturer Aviva Host Rabbit Category Antikörper Application IF, IHC, IP, WB Species Reactivity Human, Mouse, Rat Immunogen Synthesized peptide derived from human NFKappaB-p100 around the non-phosphorylation site of S865. Location: 810-890aa. This gene encodes a subunit of the transcription factor complex nuclear factor-kappa-B (NFkB). The NFkB complex is expressed in numerous cell types and functions as a central activator of genes involved in inflammation and immune function. The protei Clonality Polyclonal Concentration 1 mg/ml Molecular Weight 97 kDa NCBI Assurance ASB-OASG04979-100UL Assurance Asina Assurance Application Inflammation and immune function. The protei	Article Name	NFKB2 Antibody, Rabbit, Polyclonal
Alternative Catalog Number ASB-OASG04979-100UL Manufacturer Aviva Host Rabbit Category Antikörper Application IF, IHC, IP, WB Species Reactivity Human, Mouse, Rat Synthesized peptide derived from human NFKappaB-p100 around the non-phosphorylation site of \$865. Location: 810-890aa. This gene encodes a subunit of the transcription factor complex nuclear factor-kappa-B (NFkB). The NFkB complex is expressed in numerous cell types and functions as a central activator of genes involved in inflammation and immune function. The protei Clonality Polyclonal Concentration 1 mg/ml Molecular Weight 97 kDa NCBI Atikörper Antikörper Antikörper Antikörper Antikörper Antikörper Antikörper Antikörper Application IF, IHC, IP, WB Synthesized peptide derived from human NFKappaB-p100 around the non-phosphorylation site of \$865. Location: 810-890aa. This gene encodes a subunit of the transcription factor complex nuclear factor-kappa-B (NFkB). The NFkB complex is expressed in numerous cell types and functions as a central activator of genes involved in inflammation and immune function. The protei Clonality Polyclonal Argyl	Biozol Catalog Number	ASB-OASG04979
Manufacturer Aviva Rabbit Category Antikörper Application IF, IHC, IP, WB Species Reactivity Human, Mouse, Rat Synthesized peptide derived from human NFKappaB-p100 around the non-phosphorylation site of S865. Location: 810-890aa. This gene encodes a subunit of the transcription factor complex nuclear factor-kappa-B (NFkB). The NFkB complex is expressed in numerous cell types and functions as a central activator of genes involved in inflammation and immune function. The protei Clonality Polyclonal Concentration 1 mg/ml Molecular Weight 97 kDa NCBI 4791	Supplier Catalog Number	OASG04979
Host Rabbit Category Antikörper Application IF, IHC, IP, WB Species Reactivity Human, Mouse, Rat Synthesized peptide derived from human NFKappaB-p100 around the non-phosphorylation site of \$865. Location: 810-890aa. This gene encodes a subunit of the transcription factor complex nuclear factor-kappa-B (NFkB). The NFkB complex is expressed in numerous cell types and functions as a central activator of genes involved in inflammation and immune function. The protei Clonality Polyclonal Concentration 1 mg/ml Molecular Weight 97 kDa NCBI 4791	Alternative Catalog Number	ASB-OASG04979-100UL
Category Antikörper Application IF, IHC, IP, WB Species Reactivity Human, Mouse, Rat Immunogen Synthesized peptide derived from human NFKappaB-p100 around the non-phosphorylation site of S865. Location: 810-890aa. This gene encodes a subunit of the transcription factor complex nuclear factor-kappa-B (NFkB). The NFkB complex is expressed in numerous cell types and functions as a central activator of genes involved in inflammation and immune function. The protei Clonality Polyclonal Concentration 1 mg/ml Molecular Weight 97 kDa NCBI 4791	Manufacturer	Aviva
Application IF, IHC, IP, WB Species Reactivity Human, Mouse, Rat Immunogen Synthesized peptide derived from human NFKappaB-p100 around the non-phosphorylation site of S865. Location: 810-890aa. This gene encodes a subunit of the transcription factor complex nuclear factor-kappa-B (NFkB). The NFkB complex is expressed in numerous cell types and functions as a central activator of genes involved in inflammation and immune function. The protei Clonality Polyclonal Concentration 1 mg/ml Molecular Weight 97 kDa NCBI 4791	Host	Rabbit
Species Reactivity Human, Mouse, Rat Synthesized peptide derived from human NFKappaB-p100 around the non-phosphorylation site of S865. Location: 810-890aa. This gene encodes a subunit of the transcription factor complex nuclear factor-kappa-B (NFkB). The NFkB complex is expressed in numerous cell types and functions as a central activator of genes involved in inflammation and immune function. The protei Clonality Polyclonal Concentration 1 mg/ml Molecular Weight 97 kDa NCBI 4791	Category	Antikörper
Immunogen Synthesized peptide derived from human NFKappaB-p100 around the non-phosphorylation site of S865. Location: 810-890aa. Product Description This gene encodes a subunit of the transcription factor complex nuclear factor-kappa-B (NFkB). The NFkB complex is expressed in numerous cell types and functions as a central activator of genes involved in inflammation and immune function. The protei Clonality Polyclonal Concentration 1 mg/ml Molecular Weight 97 kDa NCBI 4791	Application	IF, IHC, IP, WB
the non-phosphorylation site of S865. Location: 810-890aa. This gene encodes a subunit of the transcription factor complex nuclear factor-kappa-B (NFkB). The NFkB complex is expressed in numerous cell types and functions as a central activator of genes involved in inflammation and immune function. The protei Clonality Polyclonal Concentration 1 mg/ml Molecular Weight 97 kDa NCBI 4791	Species Reactivity	Human, Mouse, Rat
Product Description nuclear factor-kappa-B (NFkB). The NFkB complex is expressed in numerous cell types and functions as a central activator of genes involved in inflammation and immune function. The protei Clonality Polyclonal Concentration 1 mg/ml Molecular Weight 97 kDa NCBI 4791	Immunogen	
Concentration 1 mg/ml Molecular Weight 97 kDa NCBI 4791	Product Description	nuclear factor-kappa-B (NFkB). The NFkB complex is expressed in numerous cell types and functions as a central activator of genes
Molecular Weight 97 kDa NCBI 4791	Clonality	Polyclonal
NCBI 4791	Concentration	1 mg/ml
	Molecular Weight	97 kDa
UniProt Q00653, Q9WTK5	NCBI	4791
	UniProt	Q00653, Q9WTK5

Form	Liquid. 1 mg/mL in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
------	--