

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Anti-NSE gamma (Neuron Specific Enolase, gamma) (Neuroendocrine Marker) ENO2 Monoclonal Antibody, Clone: [Clone: Clone: ENO2/1375], Mouse BOB-M02930

| Article Name               | Anti-NSE gamma (Neuron Specific Enolase, gamma) (Neuroendocrine Marker) ENO2 Monoclonal Antibody, Clone: [Clone: Clone: ENO2/1375], Mouse   |
|----------------------------|---|
| Biozol Catalog Number      | BOB-M02930  |
| Supplier Catalog Number    | M02930  |
| Alternative Catalog Number | BOB-M02930-100UG  |
| Manufacturer               | Boster Bio  |
| Host                       | Mouse   |
| Category                   | Antikörper  |
| Application                | IHC, WB   |
| Species Reactivity         | Human, Mouse, Rat   |
| Immunogen                  | A synthetic peptide of human NSE gamma (around aa416-433) (exact sequence is proprietary)   |
| Product Description        | Boster Bio Anti-NSE gamma (Neuron Specific Enolase, gamma) (Neuroendocrine Marker) ENO2 Monoclonal Antibody (Catalog M02930). Tested in WB, IHC applications. This antibody reacts with Human, Mouse, Rat |
| Clonality                  | Monoclonal  |
| Concentration              | Purified antibody with BSA and azide at 200ug/ml  |
| Clone Designation          | [Clone: Clone: ENO2/1375]   |

| Molecular Weight   | 47269 MW   |
|--------------------|--|
| UniProt            | P09104   |
| Buffer             | Prepared in 10mM PBS with 0.05% BSA & Double WITHOUT BSA & Double with 0.05% B |
| Purity             | 200ug/ml of antibody purified from Bioreactor Concentrate by Protein A/G.  |
| Form               | Liquid   |
| Application Dilute | Western Blot (1-2ug/ml)Immunohistochemistry (Formalin-fixed) (0.2-0.4ug/ml for 30 min at RT) (Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95&degC followed by cooling at RT for  |