

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **NANOG Antibody, Unconjugated, Rabbit, Polyclonal BYT-ORB1241185**

|                            |  |
|----------------------------|--|
| Article Name               | NANOG Antibody, Unconjugated, Rabbit, Polyclonal   |
| Biozol Catalog Number      | BYT-ORB1241185   |
| Supplier Catalog Number    | orb1241185   |
| Alternative Catalog Number | BYT-ORB1241185-100   |
| Manufacturer               | Biorbyt  |
| Host                       | Rabbit   |
| Category                   | Antikörper   |
| Application                | ELISA, WB  |
| Species Reactivity         | Human  |
| Immunogen                  | Antibody produced in rabbits immunized with a synthetic peptide corresponding a region of human NANOG. |
| Conjugation                | Unconjugated   |
| Product Description        | NANOG Antibody...  |
| Clonality                  | Polyclonal   |
| Concentration              | batch dependent  |
| Molecular Weight           | 33 kDa   |
| NCBI                       | <a href="#">079141</a>   |
| UniProt                    | <a href="#">Q9H9S0</a>   |

|                    |  |
|--------------------|--|
| Buffer             | Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.  |
| Form               | Liquid   |
| Target             | NANOG  |
| Application Dilute | NANOG antibody can be used for detection of NANOG by ELISA at 1:312500. NANOG antibody can be used for detection of NANOG by western blot at 0.25 µg/mL, and HRP conjugated secondary antibody should be diluted 1:50,000 - 100,000. |
| Application Notes  | NANOG antibody can be used for detection of NANOG by ELISA at 1:312500. NANOG antibody can be used for detection of NANOG by western blot at 0.25 µg/mL, and HRP conjugated secondary antibody should be diluted 1:50,000 - 100,000. |