

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Goat Anti-Mouse IgG (H+L) Rhodamine (TRITC) BWT-BS11502

Article Name Goat Anti-Mouse IgG (H+L) Rhodamine (TRITC)  Biozol Catalog Number BWT-BS11502  Supplier Catalog Number BS11502  Alternative Catalog Number BWT-BS11502-100UL  Manufacturer Bioworld Technology  Host Goat  Category Antikörper  Application FC, IF  Immunogen Mouse IgG (H+L)  Conjugation TRITC  Product Description Fluorophore: Tetramethyl rhodamine 5(and6)-isothiocyanate(TRITC) Amax=550nm,Emax=570nm Fluorophore/ProteinAbsorbanceRatio: A550/A280=0.46  Concentration 1 mg/ml  Buffer 0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, containing 1% BSA,50% glycerol  Purity chromatography using antigens coupled to agarose beads.  Application Dilute 1:50 -1:200 for most applications		
Supplier Catalog Number BS11502  Alternative Catalog Number BWT-BS11502-100UL  Manufacturer Bioworld Technology  Host Goat  Category Antikörper  Application FC, IF  Immunogen Mouse IgG (H+L)  Conjugation TRITC  Product Description Fluorophore: Tetramethyl rhodamine 5(and6)-isothiocyanate(TRITC) Amax=550nm,Emax=570nm Fluorophore/ProteinAbsorbanceRatio: A550/A280=0.46  Concentration 1 mg/ml  Buffer 0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6,containing 1% BSA,50% glycerol  Purity The antibody was isolated from antisera by immunoaffinity chromatography using antigens coupled to agarose beads.	Article Name	Goat Anti-Mouse IgG (H+L) Rhodamine (TRITC)
Alternative Catalog Number  Bioworld Technology  Host  Goat  Category  Antikörper  Application  FC, IF  Immunogen  Mouse IgG (H+L)  Conjugation  TRITC  Fluorophore: Tetramethyl rhodamine 5(and6)-isothiocyanate(TRITC)  Amax=550nm,Emax=570nm Fluorophore/ProteinAbsorbanceRatio: A550/A280=0.46  Concentration  1 mg/ml  Buffer  0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6,containing 1% BSA,50% glycerol  Purity  The antibody was isolated from antisera by immunoaffinity chromatography using antigens coupled to agarose beads.	Biozol Catalog Number	BWT-BS11502
Manufacturer  Bioworld Technology  Host  Goat  Category  Antikörper  Application  FC, IF  Immunogen  Mouse IgG (H+L)  Conjugation  TRITC  Product Description  Fluorophore: Tetramethyl rhodamine 5(and6)-isothiocyanate(TRITC)  Amax=550nm,Emax=570nm Fluorophore/ProteinAbsorbanceRatio: A550/A280=0.46  Concentration  1 mg/ml  Buffer  0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6,containing 1% BSA,50% glycerol  Purity  The antibody was isolated from antisera by immunoaffinity chromatography using antigens coupled to agarose beads.	Supplier Catalog Number	BS11502
Host Goat  Category Antikörper  Application FC, IF  Immunogen Mouse IgG (H+L)  Conjugation TRITC  Product Description Fluorophore: Tetramethyl rhodamine 5(and6)-isothiocyanate(TRITC)  Amax=550nm,Emax=570nm Fluorophore/ProteinAbsorbanceRatio: A550/A280=0.46  Concentration 1 mg/ml  Buffer 0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6,containing 1% BSA,50% glycerol  The antibody was isolated from antisera by immunoaffinity chromatography using antigens coupled to agarose beads.	Alternative Catalog Number	BWT-BS11502-100UL
Category Antikörper  Application FC, IF  Immunogen Mouse IgG (H+L)  Conjugation TRITC  Fluorophore: Tetramethyl rhodamine 5(and6)-isothiocyanate(TRITC) Amax=550nm,Emax=570nm Fluorophore/ProteinAbsorbanceRatio: A550/A280=0.46  Concentration 1 mg/ml  Buffer 0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6,containing 1% BSA,50% glycerol  Purity The antibody was isolated from antisera by immunoaffinity chromatography using antigens coupled to agarose beads.	Manufacturer	Bioworld Technology
Application FC, IF  Immunogen Mouse IgG (H+L)  Conjugation TRITC  Fluorophore: Tetramethyl rhodamine 5(and6)-isothiocyanate(TRITC) Amax=550nm,Emax=570nm Fluorophore/ProteinAbsorbanceRatio: A550/A280=0.46  Concentration 1 mg/ml  Buffer 0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6,containing 1% BSA,50% glycerol  Purity The antibody was isolated from antisera by immunoaffinity chromatography using antigens coupled to agarose beads.	Host	Goat
ImmunogenMouse IgG (H+L)ConjugationTRITCProduct DescriptionFluorophore: Tetramethyl rhodamine 5(and6)-isothiocyanate(TRITC) Amax=550nm,Emax=570nm Fluorophore/ProteinAbsorbanceRatio: A550/A280=0.46Concentration1 mg/mlBuffer0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6,containing 1% BSA,50% glycerolPurityThe antibody was isolated from antisera by immunoaffinity chromatography using antigens coupled to agarose beads.	Category	Antikörper
Conjugation  TRITC  Fluorophore: Tetramethyl rhodamine 5(and6)-isothiocyanate(TRITC)  Amax=550nm,Emax=570nm Fluorophore/ProteinAbsorbanceRatio: A550/A280=0.46  Concentration  1 mg/ml  Buffer  0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6,containing 1% BSA,50% glycerol  The antibody was isolated from antisera by immunoaffinity chromatography using antigens coupled to agarose beads.	Application	FC, IF
Fluorophore: Tetramethyl rhodamine 5(and6)-isothiocyanate(TRITC) Amax=550nm,Emax=570nm Fluorophore/ProteinAbsorbanceRatio: A550/A280=0.46  Concentration 1 mg/ml  Buffer 0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6,containing 1% BSA,50% glycerol  The antibody was isolated from antisera by immunoaffinity chromatography using antigens coupled to agarose beads.	Immunogen	Mouse IgG (H+L)
Product Description  Amax=550nm,Emax=570nm Fluorophore/ProteinAbsorbanceRatio: A550/A280=0.46  Concentration  1 mg/ml  0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6,containing 1% BSA,50% glycerol  The antibody was isolated from antisera by immunoaffinity chromatography using antigens coupled to agarose beads.	Conjugation	TRITC
Buffer  0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6,containing 1% BSA,50% glycerol  The antibody was isolated from antisera by immunoaffinity chromatography using antigens coupled to agarose beads.	Product Description	Amax=550nm,Emax=570nm Fluorophore/ProteinAbsorbanceRatio:
Purity  BSA,50% glycerol  The antibody was isolated from antisera by immunoaffinity chromatography using antigens coupled to agarose beads.	Concentration	1 mg/ml
chromatography using antigens coupled to agarose beads.	Buffer	
Application Dilute 1:50 -1:200 for most applications	Purity	
	Application Dilute	1:50 -1:200 for most applications