

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Recombinant Human TARC (rHuTARC/CCL17) BWT-PR1118

Recombinant Human TARC (rHuTARC/CCL17)
BWT-PR1118
PR1118
BWT-PR1118-5UG,BWT-PR1118-20UG,BWT-PR1118-1.0MG
Bioworld Technology
Biochemikalien
CCL17 is a novel CC chemokine recently identified using a signal sequence trap method. CCL17 cDNA encodes a highly basic 94 amino acid residue precursor protein with a 23 aa residue signal peptide that is cleaved to generate the 71 aa residue mature
8.0kDa, a single non-glycosylated polypeptide chain containing 71 amino acids.
Escherichia coli.
>97% by SDS-PAGE and HPLC analyses.
Sterile Filtered White lyophilized (freeze-dried) powder.
ARGTNVGREC CLEYFKGAIP LRKLKTWYQT SEDCSRDAIV FVTVQGRAIC SDPNNKRVKN AVKYLQSLER S
Lyophilized from a 0.2mm filtered concentrated solution in 20mM PB, pH 7.4, 150mM NaCl.

Application Notes

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportio