

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Anti-Macrophage L1 Protein S100A8 Monoclonal Antibody, Clone: [Clone: Clone: SPM281], Mouse BOB-M00413

| Article Name               | Anti-Macrophage L1 Protein S100A8 Monoclonal Antibody, Clone: [Clone: Clone: SPM281], Mouse   |
|----------------------------|---|
| Biozol Catalog Number      | BOB-M00413  |
| Supplier Catalog Number    | M00413  |
| Alternative Catalog Number | BOB-M00413-100UG  |
| Manufacturer               | Boster Bio  |
| Host                       | Mouse   |
| Category                   | Antikörper  |
| Application                | FC, IF, IHC   |
| Species Reactivity         | Bovine, Canine, Equine, Feline, Goat, Guinea pig, Human, Monkey,<br>Mouse, Porcine, Primate, Rabbit, Rat  |
| Immunogen                  | Affinity Purified monocyte membrane preparation   |
| Product Description        | Boster Bio Anti-Macrophage L1 Protein S100A8 Monoclonal Antibody (Catalog M00413). Tested in Flow Cytometry, IF, IHC applications. This antibody reacts with Human, Baboon, Monkey, Cow, Pig, Goat, Horse, Cat, Dog, Rabbit, Guinea pig, Rat, Mouse |
| Clonality                  | Monoclonal  |
| Concentration              | Purified antibody with BSA and azide at 200ug/ml  |
| Clone Designation          | [Clone: Clone: SPM281]  |
| Molecular Weight           | 10835 MW  |

| UniProt            | P05109  |
|--------------------|---|
| Buffer             | Prepared in 10mM PBS with 0.05% BSA & D.05% azide. Also available WITHOUT BSA & D.05% azide at 1.0mg/ml.  |
| Purity             | 200ug/ml of antibody purified from Bioreactor Concentrate by Protein A/G.   |
| Form               | Liquid  |
| Application Dilute | Flow Cytometry (1-2ug/million cells)Immunofluorescence (1-2ug/ml)Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 |