

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Mouse IgA Polyclonal Antibody, Cy3 Conjugated, Rabbit BSS-BS-0774R-CY3

Article Name Mouse IgA Polyclonal Antibody, Cy3 Conjugated, Rabbit Biozol Catalog Number BSS-BS-0774R-CY3 Supplier Catalog Number BSS-BS-0774R-CY3 Alternative Catalog Number BSS-BS-0774R-CY3-100 Manufacturer Bioss Host Rabbit Category Kits/Assays Application IF, WB Species Reactivity Mouse Conjugation Cy3 Mouse IgA (immunoglobulin A) is a glycosylated protein of 160 kDa and is produced as a monomer or as a J chain-linked dimer. Monomeric IgA constitutes 5-15 % of the serum immunoglobulins whereas dimeric IgA is localized to mucosa surfaces such as sal Clonality Polyclonal Concentration 1ug/ul Buffer Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Source Mouse IgA protein purified from serum		
Supplier Catalog Number BS-0774R-CY3 Alternative Catalog Number BSS-BS-0774R-CY3-100 Manufacturer Bioss Host Rabbit Category Kits/Assays Application IF, WB Species Reactivity Mouse Conjugation Cy3 Product Description Mouse IgA (immunoglobulin A) is a glycosylated protein of 160 kDa and is produced as a monomer or as a J chain-linked dimer. Monomeric IgA constitutes 5-15 % of the serum immunoglobulins whereas dimeric IgA is localized to mucosa surfaces such as sal Clonality Polyclonal Concentration 1ug/ul Buffer Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.	Article Name	Mouse IgA Polyclonal Antibody, Cy3 Conjugated, Rabbit
Alternative Catalog Number BSS-BS-0774R-CY3-100 Manufacturer Bioss Host Rabbit Category Kits/Assays Application IF, WB Species Reactivity Mouse Conjugation Cy3 Mouse IgA (immunoglobulin A) is a glycosylated protein of 160 kDa and is produced as a monomer or as a J chain-linked dimer. Monomeric IgA constitutes 5-15 % of the serum immunoglobulins whereas dimeric IgA is localized to mucosa surfaces such as sal Clonality Polyclonal Concentration 1ug/ul Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.	Biozol Catalog Number	BSS-BS-0774R-CY3
Manufacturer Bioss Rabbit Category Kits/Assays Application IF, WB Species Reactivity Mouse Conjugation Cy3 Mouse IgA (immunoglobulin A) is a glycosylated protein of 160 kDa and is produced as a monomer or as a J chain-linked dimer. Monomeric IgA constitutes 5-15 % of the serum immunoglobulins whereas dimeric IgA is localized to mucosa surfaces such as sal Clonality Polyclonal Concentration 1ug/ul Buffer Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.	Supplier Catalog Number	BS-0774R-CY3
Host Rabbit Category Kits/Assays Application IF, WB Species Reactivity Mouse Conjugation Cy3 Product Description Mouse IgA (immunoglobulin A) is a glycosylated protein of 160 kDa and is produced as a monomer or as a J chain-linked dimer. Monomeric IgA constitutes 5-15 % of the serum immunoglobulins whereas dimeric IgA is localized to mucosa surfaces such as sal Clonality Polyclonal Concentration 1ug/ul Buffer Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.	Alternative Catalog Number	BSS-BS-0774R-CY3-100
Category Kits/Assays Application IF, WB Species Reactivity Mouse Conjugation Cy3 Mouse IgA (immunoglobulin A) is a glycosylated protein of 160 kDa and is produced as a monomer or as a J chain-linked dimer. Monomeric IgA constitutes 5-15 % of the serum immunoglobulins whereas dimeric IgA is localized to mucosa surfaces such as sal Clonality Polyclonal Concentration 1ug/ul Buffer Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.	Manufacturer	Bioss
Application IF, WB Species Reactivity Mouse Conjugation Cy3 Mouse IgA (immunoglobulin A) is a glycosylated protein of 160 kDa and is produced as a monomer or as a J chain-linked dimer. Monomeric IgA constitutes 5-15 % of the serum immunoglobulins whereas dimeric IgA is localized to mucosa surfaces such as sal Clonality Polyclonal Concentration 1ug/ul Buffer Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.	Host	Rabbit
Species Reactivity Conjugation Cy3 Mouse IgA (immunoglobulin A) is a glycosylated protein of 160 kDa and is produced as a monomer or as a J chain-linked dimer. Monomeric IgA constitutes 5-15 % of the serum immunoglobulins whereas dimeric IgA is localized to mucosa surfaces such as sal Clonality Polyclonal Concentration 1ug/ul Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.	Category	Kits/Assays
Conjugation Cy3 Mouse IgA (immunoglobulin A) is a glycosylated protein of 160 kDa and is produced as a monomer or as a J chain-linked dimer. Monomeric IgA constitutes 5-15 % of the serum immunoglobulins whereas dimeric IgA is localized to mucosa surfaces such as sal Clonality Polyclonal Concentration 1ug/ul Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.	Application	IF, WB
Product Description Mouse IgA (immunoglobulin A) is a glycosylated protein of 160 kDa and is produced as a monomer or as a J chain-linked dimer. Monomeric IgA constitutes 5-15 % of the serum immunoglobulins whereas dimeric IgA is localized to mucosa surfaces such as sal Clonality Polyclonal Concentration 1ug/ul Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.	Species Reactivity	Mouse
Product Description and is produced as a monomer or as a J chain-linked dimer. Monomeric IgA constitutes 5-15 % of the serum immunoglobulins whereas dimeric IgA is localized to mucosa surfaces such as sal Clonality Polyclonal Concentration 1ug/ul Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.	Conjugation	СуЗ
Concentration 1ug/ul Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.	Product Description	and is produced as a monomer or as a J chain-linked dimer. Monomeric IgA constitutes 5-15 % of the serum immunoglobulins
Buffer Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.	Clonality	Polyclonal
BSA, 0.02% Proclin300 and 50% Glycerol.	Concentration	1ug/ul
Source Mouse IgA protein purified from serum	Buffer	
	Source	Mouse IgA protein purified from serum
Purity affinity Purified by Protein A.	Purity	affinity Purified by Protein A.

Target	Mouse IgA
Application Dilute	WB(1:300-5000), IF(IHC-P)(1:50-200), IF(IHC-F)(1:50-200)