

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Alexa Fluor 647-conjugated AffiniPure(TM) Sheep Anti-Mouse IgG (H+L) (min X Hu,Bov,Hrs Sr Prot) JIM-515-605-062

| Article Name | Alexa Fluor 647-conjugated AffiniPure(TM) Sheep Anti-Mouse IgG (H+L) (min X Hu,Bov,Hrs Sr Prot) |
|----------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Biozol Catalog Number | JIM-515-605-062 |
| Supplier Catalog Number | 515-605-062 |
| Alternative Catalog Number | JIM-515-605-062 |
| Manufacturer | Jackson ImmunoResearch |
| Host | Sheep |
| Category | Antikörper |
| Species Reactivity | Mouse |
| Conjugation | Alexa Fluor® 647 |
| Clonality | Polyclonal |
| Buffer | Buffer: 0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6. Stabilizer: 15 mg/ml Bovine Serum Albumin (IgG-Free, Protease-Free). Preservative: 0.05% Sodium Azide |
| Purity | AffiniPure |
| Form | Freeze-dried solid |
| Storage | Storage and Rehydration: Store freeze-dried solid at 2-8°C. Rehydrate with the indicated volume of dH2O and centrifuge if not clear. Prepare working dilution on day of use. Product is stable for about 6 weeks at 2-8°C as an undiluted liquid. Extended |

| Antibody Type | Secondary Antibody |
|--------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Application Dilute | 1:100 - 1:800 for most applications Dilution factors are presented in the form of a range because the optimal dilution is a function of many factors, such as antigen density, permeability, etc. The actual dilution used must be determined empirically. |