

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Rhodamine (TRITC)-conjugated AffiniPure(TM) Donkey Anti-Goat IgG (H+L) (min X Ck,GP,Sy Hms,Hrs,Hu,Ms,Rb,Rat Sr Prot) JIM-705-025-147

| | |
|----------------------------|---|
| Article Name | Rhodamine (TRITC)-conjugated AffiniPure(TM) Donkey Anti-Goat IgG (H+L) (min X Ck,GP,Sy Hms,Hrs,Hu,Ms,Rb,Rat Sr Prot) |
| Biozol Catalog Number | JIM-705-025-147 |
| Supplier Catalog Number | 705-025-147 |
| Alternative Catalog Number | JIM-705-025-147 |
| Manufacturer | Jackson ImmunoResearch |
| Host | Donkey |
| Category | Antikörper |
| Species Reactivity | Goat |
| Conjugation | TRITC |
| Clonality | Polyclonal |
| Buffer | Buffer: 0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6. Stabilizer: 15 mg/ml Bovine Serum Albumin (IgG-Free, Protease-Free). Preservative: 0.05% Sodium Azide |
| Purity | AffiniPure |
| Form | Freeze-dried solid |
| Storage | Storage and Rehydration: Store freeze-dried solid at 2-8°C. Rehydrate with the indicated volume of dH2O and centrifuge if not clear. Prepare working dilution on day of use. Product is stable for about 6 weeks at 2-8°C as an undiluted liquid. Extended... |

| | |
|--------------------|---|
| Antibody Type | Secondary Antibody |
| Application Dilute | 1:50 - 1:200 for most applications Dilution factors are presented in the form of a range because the optimal dilution is a function of many factors, such as antigen density, permeability, etc. The actual dilution used must be... |