

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Rhodamine Red-X-conjugated AffiniPure(TM) F(ab)2 Fragment Donkey Anti-Human IgG (H+L) (min X Bov,Ck,Gt,GP,Sy Hms,Hrs,Ms,Rb,Rat,Shp Sr Prot)

JIM-709-296-149

| Article Name | Rhodamine Red-X-conjugated AffiniPure(TM) F(ab)2 Fragment Donkey Anti-Human IgG (H+L) (min X Bov,Ck,Gt,GP,Sy Hms,Hrs,Ms,Rb,Rat,Shp Sr Prot) |
|----------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------|
| Biozol Catalog Number | JIM-709-296-149 |
| Supplier Catalog Number | 709-296-149 |
| Alternative Catalog Number | JIM-709-296-149 |
| Manufacturer | Jackson ImmunoResearch |
| Host | Donkey |
| Category | Antikörper |
| Species Reactivity | Human |
| Conjugation | Rhodamine Red™-X |
| Clonality | Polyclonal |
| Buffer | Buffer: 0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6. Stabilizer: 15 mg/ml Bovine Serum Albumin (IgG-Free, Protease-Free). Preservative: 0.05% Sodium Azide |
| Purity | AffiniPure |
| Form | Freeze-dried solid |

| Storage | Storage and Rehydration: Store freeze-dried solid at 2-8°C. Rehydrate with the indicated volume of dH2O and centrifuge if not clear. Prepare working dilution on day of use. Product is stable for about 6 weeks at 2-8°C as an undiluted liquid. Extended |
|--------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Antibody Type | Secondary Antibody |
| Application Dilute | _x000D_ 1:50 - 1:200 for most applications_x000Dx000D_ Dilution factors are presented in the form of a range because the optimal dilution is a function of many factors, such as antigen density, permeability, etc. The actual dilution used must be |