

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Sheep anti Bovine IgG2 (subclass specific), Clone: [Polyclonal], Monoclonal NMB-SHAB/IGG2

| | |
|----------------------------|--|
| Article Name | Sheep anti Bovine IgG2 (subclass specific), Clone: [Polyclonal], Monoclonal |
| Biozol Catalog Number | NMB-SHAB/IGG2 |
| Supplier Catalog Number | ShAB/IgG2 |
| Alternative Catalog Number | NMB-SHAB/IGG2 |
| Manufacturer | NordicMubio |
| Host | Sheep |
| Category | Antikörper |
| Product Description | The reactivity of the antiserum is directed to the subclass specific determinants on the Fc subunit of IgG1. It does not react with other IgG2, IgG/Fab fragments, IgM and IgA or any non-Ig protein in bovine serum, as tested by immunoelectrophoresis a... |
| Clonality | Monoclonal |
| Clone Designation | [Polyclonal] |
| Buffer | Delipidated, heat inactivated, lyophilized, stable whole antiserum. No preservative added, as it may interfere with the antibody activity. Total protein and IgG concentrations in the antiserum are comparable to those in pooled sheep serum. No foreign pro |
| Source | Pools of purified normal IgG1 isolated from pooled bovine serum. Freund's complete adjuvant is used in the first step of the immunization procedure. |

| | |
|-------------------|--|
| Formula | Delipidated, heat inactivated, lyophilized, stable whole antiserum. No preservative added, as it may interfere with the antibody activity. Total protein and IgG concentrations in the antiserum are comparable to those in pooled sheep serum. No foreign pro |
| Application Notes | Precipitation assays. In immunoelectrophoresis use 2 µl or equivalent against 120 µl antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 µl antiserum in a 3 mm diameter centre well and 2 µl serum samples (neat and |