

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Sheep anti Dog IgG (Fc specific), Clone: [Polyclonal], Monoclonal NMB-SHAD/IGG(FC)

| | |
|----------------------------|--|
| Article Name | Sheep anti Dog IgG (Fc specific), Clone: [Polyclonal], Monoclonal |
| Biozol Catalog Number | NMB-SHAD/IGG(FC) |
| Supplier Catalog Number | ShAD/IgG(Fc) |
| Alternative Catalog Number | NMB-SHAD/IGG(FC) |
| Manufacturer | NordicMubio |
| Host | Sheep |
| Category | Antikörper |
| Application | ELISA, ICC |
| Species Reactivity | Canine |
| Product Description | The reactivity of the antiserum is restricted to the Fc part of the IgG molecule. In immunoelectrophoresis and radial immunodiffusion, using various antiserum concentrations against normal dog serum a single precipitin line is obtained which shows a ... |
| Clonality | Monoclonal |
| Clone Designation | [Polyclonal] |
| Buffer | Delipidated, heat inactivated, lyophilized, stable whole antiserum. 0.01% sodium azide. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal sheep serum. No foreign proteins added. Reconstitute the lyophilized an |
| Source | Highly purified normal IgG isolated from pooled dog serum was used for immunization of the sheep. Freund's complete adjuvant is used in the first step of the immunization procedure. |

| | |
|-------------------|---|
| Formula | Delipidated, heat inactivated, lyophilized, stable whole antiserum. 0.01% sodium azide. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal sheep serum. No foreign proteins added. |
| Application Notes | Precipitation assays. In immunoelectrophoresis use 2 μ l serum or equivalent against 120 μ l antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 μ l antiserum in 3 mm diameter center well and 2 μ l serum samples (neat |