

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Sheep anti Horse IgG (heavy and light chains), Clone: [Polyclonal], Monoclonal NMB-SHAHO/IGG(H+L)

| | |
|----------------------------|--|
| Article Name | Sheep anti Horse IgG (heavy and light chains), Clone: [Polyclonal], Monoclonal |
| Biozol Catalog Number | NMB-SHAHO/IGG(H+L) |
| Supplier Catalog Number | ShAHo/IgG(H+L) |
| Alternative Catalog Number | NMB-SHAHO/IGG(H+L) |
| Manufacturer | NordicMubio |
| Host | Sheep |
| Category | Antikörper |
| Product Description | The reactivity of the antiserum is directed to the Fc and Fab subunits of both subclasses IgG1 and IgG2. Reactivity with other subclasses of horse IgG has not been tested. The antiserum reacts with both Fc and Fab portions of polyclonal IgG. It inclu... |
| Clonality | Monoclonal |
| Clone Designation | [Polyclonal] |
| Buffer | Delipidated, heat inactivated lyophilized stable whole serum. No preservative added, as it may interfere with the antibody activity. No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled sheep se |
| Source | Purified normal IgG isolated from pooled horse serum. Freund's complete adjuvant is used in the first step of the immunization procedure. |

| | |
|-------------------|--|
| Formula | Delipidated, heat inactivated lyophilized stable whole serum. No preservative added, as it may interfere with the antibody activity. No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled sheep se |
| Application Notes | Precipitation assays. In immunoelectrophoresis use 2 µl or equivalent against 120 µl antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 µl antiserum in a 3 mm diameter centre well and 2 µl serum samples (neat and |